
Pairwise Cluster Comparison for Learning Latent Variable Models

Nuaman Asbeh

Industrial Engineering and Management
Ben-Gurion University of the Negev
Beer-Sheva, 84105, Israel

Boaz Lerner

Industrial Engineering and Management
Ben-Gurion University of the Negev
Beer-Sheva, 84105, Israel

Abstract

Identification of latent variables that govern a problem and the relationships among them, given measurements in the observed world, are important for causal discovery. This identification can be accomplished by analyzing the constraints imposed by the latents in the measurements. We introduce the concept of *pairwise cluster comparison* (PCC) to identify causal relationships from clusters of data points and provide a two-stage algorithm called learning PCC (LPCC) that learns a latent variable model (LVM) using PCC. LPCC learns exogenous latents and latent colliders before also learning latent non-colliders, all together with their observed descendants, by using pairwise comparisons between data clusters in the measurement space that explain latent causes. If the true graph has no serial connections, LPCC returns the true graph, and if the true graph has at least a single serial connection, LPCC returns a pattern of the true graph. LPCC's most important advantage is that it is not limited to linear or latent-tree models and makes only mild assumptions about the distribution. The code and evaluation results are freely available (<http://www.ee.bgu.ac.il/~boaz/software.html>), and too technical details and all proofs are in a supplementary material file.

1. INTRODUCTION & RELATED WORK

Latent variable models (LVMs) represent latent variables and the causal relationships among them to explain observed variables that have been measured in the domain. LVMs reduce dimensionality by aggregating (many) observed variables into a few latent variables, each of which represents a “concept” explaining some aspects of the domain that can be interpreted from the data.

Learning an LVM exploits values of the measured variables as manifested in the data to make an inference about the causal relationships among the latent variables and to predict the value of these variables. Although statistical methods for learning an LVM, such as factor analysis, effectively reduce dimensionality and may fit the data reasonably well, the resulting models might not have any correspondence to real causal mechanisms (Silva et al., 2006). On the other hand, the focus of learning Bayesian networks (BNs) is on causal relations among observed variables, whereas the detection of latent variables and the interrelations among themselves and with the observed variables has received little attention. Learning an LVM using Inductive Causation* (IC*) (Pearl and Verma, 1991) and Fast Causal Inference (FCI) (Spirtes et al., 2000) returns partial ancestral graphs, which indicate for each link whether it is a (potential) manifestation of a hidden common cause for the two linked variables. The structural EM algorithm (Friedman, 1998) learns a structure using a fixed set of previously given latents. By searching for “structural signatures” of latents, the FindHidden algorithm (Elidan et al., 2000) detects substructures that suggest the presence of latents in the form of dense subnetworks. Elidan and Friedman (2001) give a fast algorithm for determining the cardinality of latent variables introduced this way. However, Silva et al. (2006) suspected that FindHidden cannot always find a pure measurement sub-model, which is a flaw in causal analysis. Also, the recovery of latent trees of binary and Gaussian variables has been suggested (Pearl, 2000). Hierarchical latent class (HLC) models, which are rooted trees where the leaf nodes are observed while all other nodes are latent, were proposed for clustering (Zhang, 2004). Two greedy algorithms are suggested (Harmeling and Williams, 2011) to expedite learning the structure of a binary HLC and the cardinalities of the latents. The BIN-G algorithm determines both the structure of the tree and the cardinality of the latent variables in a bottom-up fashion. The BIN-A algorithm first determines the tree structure using agglomerative hierarchical clustering and then determines the cardinality of the latent variables in the same manner as the BIN-G algorithm.

Latent-tree models are also used to speed approximate inference in BN, trading the approximation accuracy with inferential complexity (Wang et al., 2008).

Models in which multiple latents may have multiple indicators (observed children), also known as multiple indicator models (MIMs) (Bartholomew et al., 2002; Spirtes, 2013), are a very important subclass of structural equation models (SEM), which are widely used, together with BNs, to analyze causal relations (Shimizu et al., 2011). For these models, and others that are not tree-constrained, most of the mentioned algorithms may lead to unsatisfactory results. This is one of the most difficult problems in machine learning and statistics since, in general, a joint distribution can be generated by an infinite number of different LVMs. However, an algorithm that fills the gap between learning latent-tree models and learning MIMs is BuildPureClusters (BPC; Silva et al., 2006). BPC searches for the set (an equivalence class) of MIMs that best matches the set of vanishing tetrad differences (Scheines et al., 1995), but is limited to linear models (Spirtes, 2013).

In this study, we make another attempt in this direction and target the goal of Silva et al. (2006), but concentrate on the discrete case, rather than on the continuous case dealt by BPC, using ideas and principles of clustering. Interestingly, the same difficulty in learning MIMs is also faced in the domain of unsupervised learning that confronts similar questions such as: (1) How many clusters are there in the observed data? and (2) Which classes do the clusters really represent? Due to this similarity, we propose a concept and an algorithm that link learning an LVM and clustering analysis. Using the *pairwise cluster comparison* (PCC) concept, we compare pairwise clusters of data points representing instantiations of the observed variables to identify those pairs of clusters that exhibit major changes in the observed variables due to changes in their ancestor latent variables. Changes in a latent variable that are manifested in changes in the observed variables reveal this latent variable and its causal paths of influence in the domain. Using the *learning PCC* (LPCC) algorithm, we learn an LVM. We identify PCCs and use them to learn latent variables – exogenous and endogenous (the latter may be either colliders or non-colliders) – and their causal interrelationships as well as their children (latent variables and observed variables) and causal paths from latent variables to observed variables.

2. LVM LEARNING

The goal of our study is to reconstruct an LVM from i.i.d. data sampled from the observed variables in an unknown model. First, we present the assumptions that LPCC makes and the constraints it applies on LVM and compare them to those required by other state-of-the-art methods.

Assumption 1 The underlying model is a Bayesian network, $BN = \langle G, \Theta \rangle$, encoding a discrete joint probability distribution P for a set of random variables $V = L \cup O$, where $G = \langle V, E \rangle$ is a directed acyclic graph (DAG) whose nodes V correspond to latents L and observed variables O , and E is the set of edges between nodes in G . Θ is the set of parameters, i.e., the conditional probabilities of variables in V given their parents.

Assumption 2 No observed variable in O is an ancestor of any latent variable in L . This property is called the measurement assumption (Spirtes et al., 2000).

Before we present additional assumptions about the learned LVM, we need Definitions 1–4 (following Silva et al., 2006), which are specific to LVM:

Definition 1 A model satisfying Assumptions 1 and 2 is a latent variable model.

Definition 2 Given an LVM G with a variable set V , the subgraph containing all variables in V and all and only those edges directed into variables in O is called the *measurement model* of G .

Definition 3 Given an LVM G , the subgraph containing all and only G 's latent nodes and their respective edges is called the *structural model* of G .

When each model variable is a linear function of its parents in the graph plus an additive error term of positive finite variance, the latent variable model is linear; this is also known as SEM. In this study, we dispense with the linearity assumption and apply the above concepts to learn not necessarily linear MIMs or latent-tree models. Our suggested concept, LPCC, is not limited by the linearity assumption and learns a model as long as it is MIM. In addition, we are interested in discrete LVMs. Another important definition we need is:

Definition 4 A *pure measurement model* is a measurement model in which each observed variable has only one latent parent and no observed parent.

Assumption 3 The measurement model of G is pure.

As a principled way of testing conditional independence among latents, Silva et al. (2006) focus on MIMs, which are pure measurement models. Practically, these models have a smaller equivalence class of the latent structure than that of non-pure models and thus are easier to learn unambiguously.

LPCC does not assume that the true measurement model is linear (which is a parametric assumption that e.g., BPC makes), but rather assumes that the model is pure (a structural assumption). When the true causal model is pure, LPCC will identify it correctly (or find its pattern). When

it is not pure, LPCC – similarly to BPC (Silva et al., 2006) – will learn a pure sub-model of the true model using two indicators for each latent (compared to three indicators per latent that are required by BPC). In addition, LPCC assumes that:

Assumption 4 The true model \mathbf{G} is MIM, in which each latent has at least two observed children and may have latent parents.

Causal structure discovery – learning the number of latent variables in the model, their interconnections and connections to the observed variables, as well as the interconnections among the observed variables – is very difficult and thus requires making some assumptions about the problem. Particularly, MIMs, in which multiple observed variables are assumed to be affected by latent variables and perhaps by each other (Spirtes, 2013), are reasonable models but have attracted scant attention in the machine-learning community. By assuming that the true model manifests local influence of each latent variable on at least a small number of observed variables, Silva et al. (2006) showed that learning the complete Markov equivalence class of MIM is feasible. Similar to Silva et al. (2006), we assume that the true model is MIM.

3. PRELIMINARIES TO LPCC

Figure 1 sketches a range of MIMs, which all exhibit pure measurement models, presenting a range of challenges to an LVM learning algorithm. Compared to G_1 , which is a basic MIM of two unconnected latents, G_2 shows a structural model that is characterized by a latent collider. Note that such an LVM cannot be learned by latent-tree algorithms such as in Zhang (2004). G_3 and G_4 demonstrate serial and diverging structural models, respectively, that together with G_2 cover the three basic structural models. G_5 and G_6 manifest more complex structural models comprising a latent collider and a combination of serial and diverging connections.

In Section 3.1, we show that learning pairwise cluster comparison relies on understanding the influence of the exogenous latent variables on the observed variables in the LVM. This influence is divided into major and minor effects that are introduced and explained in Section 3.2. In Section 3.3, we link this structural influence to data clustering and introduce the pairwise cluster comparison concept for learning an LVM.

3.1 THE INFLUENCE OF EXOGENOUS LATENTS

We distinguish between observed (\mathbf{O}) and latent (\mathbf{L}) variables and between exogenous (\mathbf{EX}) and endogenous (\mathbf{EN}) variables. \mathbf{EX} have zero in-degree, are autonomous, and unaffected by the values of the other variables (e.g., L_1 in all graphs but G_4 in Figure 1), whereas \mathbf{EN} are all non-exogenous variables in \mathbf{G} (e.g., L_2 in all graphs but G_1 and G_4 , and X_1 in all graphs in Figure 1). We identify three

types of variables: (1) Exogenous latents, $\mathbf{EX} \subset (\mathbf{L} \cap \mathbf{NC})$ [all exogenous variables are latent non-colliders (\mathbf{NC})]; (2) Endogenous latents, $\mathbf{EL} \subset (\mathbf{L} \cap \mathbf{EN})$, which are divided into latent colliders $\mathbf{C} \subset \mathbf{EL}$ (e.g., L_2 in G_2 and G_5 ; note that all latent colliders are endogenous) and latent non-colliders (in serial and diverging connections) $\mathbf{S} \subset (\mathbf{EL} \cap \mathbf{NC})$ (e.g., L_3 in G_3 , G_4 , and G_6), thus $\mathbf{NC} = (\mathbf{EX} \cup \mathbf{S})$; and (3) Observed variables, $\mathbf{O} \subset \mathbf{EN}$, which are always exogenous and childless, that are divided into children of exogenous latents $\mathbf{OEX} \subset \mathbf{O}$ (e.g., X_1 and X_9 in G_2), children of latent colliders $\mathbf{OC} \subset \mathbf{O}$ (e.g., X_4 , X_5 , and X_6 in G_2), and children of endogenous latent non-colliders $\mathbf{OS} \subset \mathbf{O}$ (e.g., X_4 , X_5 , and X_6 in G_3). We denote value configurations of \mathbf{EX} , \mathbf{EN} (when we do not know whether the endogenous variables are latent or observed), \mathbf{EL} , \mathbf{C} , \mathbf{NC} (when we do not know whether the non-collider variables are exogenous or endogenous), \mathbf{S} , \mathbf{O} , \mathbf{OEX} , \mathbf{OC} , and \mathbf{OS} by \mathbf{ex} , \mathbf{en} , \mathbf{el} , \mathbf{c} , \mathbf{nc} , \mathbf{s} , \mathbf{o} , \mathbf{oex} , \mathbf{oc} , and \mathbf{os} , respectively.

In this paper, we claim and demonstrate that the influence of exogenous (latent) variables on observed variables is fundamental to learning an LVM and introduce LPCC that identifies and exploits this influence on learning an MIM. In this section, we prove that changes in values of the observed variables are due to changes in values of the exogenous variables and thus the identification of the former indicates the existence of the latter. To do that, we analyze the propagation of influence along paths connecting both variables, remembering that the paths may contain latent colliders and latent non-colliders. First, however, we should analyze paths among the latents and only then paths ending in their sinks (i.e., the observed variables).

While BPC (Silva et al., 2006) needs to make a parametric assumption about the linearity of the model, LPCC assumes on the model structure (Assumption 3 above and Assumption 5 below). This is also the approach of latent-tree algorithms (Zhang, 2004; Harmeling and Williams, 2011; Wang et al., 2008) that restrict the learned structure to a tree (note that LPCC is not limited to a tree because it allows latent variables to be colliders). This shows a tradeoff between the structural and parametric assumptions that an algorithm for learning an LVM usually has to make; the fewer parametric assumptions the algorithm makes, the more structural assumptions it has to make and vice versa.

Assumption 5 A latent collider does not have any latent descendants.

To distinguish between latent colliders and latent non-colliders, their observed children, and their connectivity patterns to their exogenous variables, we use Lemma 1. Latent colliders and their observed children are connected to several exogenous variables via several directed paths, whereas latent non-colliders and their observed children are connected only to a single exogenous variable via a

single directed path [the necessary Definitions 5-7 are in a supplementary material file (SMF)]. Use of these different connectivity patterns – from exogenous latents through endogenous latents (both colliders and non-colliders) to observed variables – simplifies the analysis of the influence of latents on observed variables. Proposition 1 quantifies these patterns through the joint probability distribution.

Lemma 1 1) Each latent non-collider NC_t has only one exogenous latent ancestor EX_{NC_t} , and there is only one directed path T_{NC_t} from EX_{NC_t} (source) to NC_t (sink). Note that we use the notation NC_t , rather than S_t , since the Lemma applies to both exogenous and endogenous latent non-colliders. 2) Each latent collider C_j is connected to a set of exogenous latent ancestors EX_{C_j} via a set of directed paths T_{C_j} from EX_{C_j} (sources) to C_j (sink).

Proposition 1 The joint probability over \mathbf{V} due to value assignment \mathbf{ex} to exogenous set \mathbf{EX} is determined only by this assignment and the BN conditional probabilities.

3.2 MAJOR AND MINOR EFFECTS & VALUES

So far, we have analyzed the structural influences (path of hierarchies) of the latents on the observed variables. In this section, we complement this analysis with the parametric influences, which we divide into major and minor effects. First, we define local influence on a single EN of its direct parents. Second, we use local influences and the BN Markov property to generalize the influence of \mathbf{EX} on \mathbf{EN} . Third, exploiting the connectivity between the exogenous ancestors and their endogenous descendants, as described by Lemma 1, we focus on the influence of a specific (partial) set of exogenous variables on the values of their endogenous descendants. Analysis of the influence of all configurations \mathbf{ex} on all \mathbf{ens} and that of the configurations of specific exogenous ancestors in these \mathbf{ex} on their endogenous descendants enable learning the structure and parameters of the model and causal discovery. In Section 3.3, we show how these concepts can be exploited to learn an LVM from data clustering.

Definition 8 (definition and assumption numbers consider all those in the SMF) A *local effect* on an endogenous variable EN is the influence of a configuration of EN 's direct latent parents on any of EN 's values. Following, we define a major local effect as the largest local effect on EN and a minor local effect as any non-major local effect on EN .

A *major local effect* on EN_i is identified by the maximal conditional probability of a specific en_i given a configuration \mathbf{pa}_i of EN_i 's latent parents \mathbf{Pa}_i , such that $MAE_{EN_i}(\mathbf{pa}_i) = \max_{en_i'}\{P(EN_i = en_i' | \mathbf{Pa}_i = \mathbf{pa}_i)\}$. All probabilities of other values of EN_i conditioned on \mathbf{pa}_i that are smaller than $MAE_{EN_i}(\mathbf{pa}_i)$ identify the *minor local effect* set, $MIES_{EN_i}(\mathbf{pa}_i)$. Similarly, the

major local value is the en_i corresponding to $MAE_{EN_i}(\mathbf{pa}_i)$, i.e., the most probable value of EN_i due to \mathbf{pa}_i , $MAV_{EN_i}(\mathbf{pa}_i) = \operatorname{argmax}_{en_i'}\{P(EN_i = en_i' | \mathbf{Pa}_i = \mathbf{pa}_i)\}$. A *minor local value* is an en_i corresponding to a minor local effect, and $MIVS_{EN_i}(\mathbf{pa}_i)$ is the set of all minor values that correspond to $MIES_{EN_i}(\mathbf{pa}_i)$. When EN_i is an observed variable or an endogenous latent non-collider, and thus has only a single parent Pa_i , the configuration \mathbf{pa}_i is actually the value pa_i of Pa_i .

We: 1) assume (Assumption 6) that for every endogenous variable EN_i in \mathbf{G} and every configuration \mathbf{pa}'_i of EN_i 's parents \mathbf{Pa}_i , there exists a certain value en'_i of EN_i , s.t. $P(EN_i = en'_i | \mathbf{Pa}_i = \mathbf{pa}'_i) > P(EN_i = en''_i | \mathbf{Pa}_i = \mathbf{pa}'_i)$ for every other value en''_i of EN_i (Pearl, 1988); 2) prove (Proposition 2 in the SMF) that $MAV_{EN_i}(\mathbf{pa}'_i)$ is certain given a certain \mathbf{pa}'_i ; and 3) assume (Assumption 7) that a) for every EN_i that is an observed variable or an endogenous latent non-collider and for every two values pa'_i and pa''_i of Pa_i , $MAV_{EN_i}(pa'_i) \neq MAV_{EN_i}(pa''_i)$, and b) for every C_j that is a latent collider and for every $Pa_j \in \mathbf{Pa}_j$, there are at least two configurations \mathbf{pa}'_j and \mathbf{pa}''_j of \mathbf{Pa}_j in which only the value of Pa_j is different and $MAV_{C_j}(\mathbf{pa}'_j) \neq MAV_{C_j}(\mathbf{pa}''_j)$.

By aggregation over all local influences, we can now generalize these concepts through the BN parameters and Markov property from local influences on specific endogenous variables to influence on all endogenous variables in the graph.

Then, we define (Definition 9): 1) an *effect* on \mathbf{EN} as the influence of a configuration \mathbf{ex} of \mathbf{EX} on \mathbf{EN} ; 2) a *major effect* (MAE) as the largest effect of \mathbf{ex} on \mathbf{EN} and a *minor effect* (MIE) as any non-MAE effect of \mathbf{ex} on \mathbf{EN} ; and 3) a *major value configuration* (MAV) as the configuration \mathbf{en} of \mathbf{EN} corresponding to MAE (i.e., the most probable \mathbf{en} due to \mathbf{ex}), and a *minor value configuration* is a configuration \mathbf{en} corresponding to any MIE.

A configuration \mathbf{en} of \mathbf{EN} in which each variable in \mathbf{EN} takes on the major local value is major or a MAV. Any effect in which at least one EN takes on a minor local effect is minor, and any configuration in which at least one EN takes on a minor local value is minor. We denote the set of all minor effects for \mathbf{ex} with $MIES(\mathbf{ex})$ (with correspondence to $MIES_{EN_i}$) and the set of all minor configurations with $MIVS(\mathbf{ex})$ (with correspondence to $MIVS_{EN_i}$).

Now, we are interested in representing the influence on a subset of the endogenous variables of the subset of the exogenous variables that impact these endogenous variables. This partial representation of MAE will enable LPCC to recover the relationships between exogenous ancestors and only the descendants that are

affected by these exogenous variables. To achieve this, we first extend the concept of effect to the concept of partial effect of specific exogenous variables and then quantify it.

We separately analyze the partial effect (Definition 10 in the SMF) of each exogenous variable on each observed variable for which the exogenous is its ancestor and all the latent variables along the path connecting these two and distinguish between two cases: (1) Observed descendants in **OEX** and **OS** that are, respectively, children of exogenous latents and children of latent non-colliders that are linked to their exogenous ancestors, each via a single directed path (e.g., the partial effect of a value of L3 in G5 (Figure 1) on {L4, L5, X13}); and (2) Observed descendants in **OC** that are children of latent colliders and linked to their exogenous ancestors via a set of directed paths through their latent collider parents (e.g., the partial effect of a configuration of $\mathbf{EX}_{C_k} = \mathbf{EX}_{L_4} = \{L1, L5\}$ in G6 (Figure 1) on {L2, L3, L4, X11}).

In the SMF, we provide detailed descriptions for these partial effects and values for observed children of latent non-colliders (*Case 1*) and observed children of latent colliders (*Case 2*) and formalize their properties (Propositions 3–7) to set the stage for Lemma 2.

Lemma 2 1) A latent non-collider NC_v and its observed child ONC_v , both descendants of an exogenous variable EX_{NC_v} , change their values in any two major configurations if and only if EX_{NC_v} has changed its value in the corresponding two configurations of \mathbf{EX} .; and 2) A latent collider C_k and its observed child OC_k , both descendants of a set of exogenous variables \mathbf{EX}_{C_k} , change their values in any two major configurations, only if at least one of the exogenous variables in \mathbf{EX}_{C_k} has changed its value in the corresponding two configurations of \mathbf{EX} .

3.3 PCC BY DATA CLUSTERING

Practically, we use observational data that were generated from an unknown LVM and measured over the observed variables. Proposition 1 showed us that each configuration of observed variables (which is part of a configuration of the endogenous variables) and their joint probability is a result of an assignment of a configuration \mathbf{ex} to the exogenous variables \mathbf{EX} . Therefore, we 1) define (Definition 11) an *observed value configuration*, *observed major value configuration*, and *observed minor value configuration* due to \mathbf{ex} that are the parts in \mathbf{en} , MAV, and a minor value configuration, respectively, that correspond to the observed variables; and 2) prove (Propositions 8 and 9 in the SMF) that a) there is only a single observed major value configuration to each exogenous configuration \mathbf{ex} ; and b) there are different observed major value configurations to different exogenous configurations \mathbf{exs} .

Due to the probabilistic nature of BN, each observed value configuration due to \mathbf{ex} may be represented by several data points. Clustering these data points may produce several clusters for each \mathbf{ex} , and each cluster corresponds to another observed value configuration. Based on Propositions 8 and 9, one and only one of the clusters corresponds to each of the observed major value configurations, whereas the other clusters correspond to observed minor value configurations. We distinguish between these clusters using the following definition:

Definition 12 The single cluster that corresponds to the observed major value configuration, and thus also represents the major effect $MAE(\mathbf{ex})$ due to configuration \mathbf{ex} , is the *major cluster* for \mathbf{ex} , and all the clusters that correspond to the observed minor value configurations due to minor effects in $MIES(\mathbf{ex})$ are *minor clusters*.

To resolve between different types of minor effects/clusters, we make two definitions.

Definition 13 A k-order minor effect is a minor effect in which exactly k endogenous variables in EN correspond to minor local effects. An \mathbf{en} corresponding to a k-order minor effect is a k-order minor value configuration.

Definition 14 Minor clusters that correspond to k-order minor effects are k-order minor clusters.

Based on Proposition 9 and Definition 12, the set of all major clusters (corresponding to all observed major value configurations) reflects the effect of all possible \mathbf{exs} , and thus the number of major clusters is expected to be equal to the number of \mathbf{EX} configurations. Therefore, the identification of all major clusters is a key to the discovery of exogenous variables and their causal interrelations. For this purpose, we introduce the concept of pairwise cluster comparison (PCC). PCC measures the differences between two clusters; each represents the response of LVM to another \mathbf{ex} .

Definition 15 Pairwise cluster comparison is a procedure by which pairs of clusters are compared, for example, through a comparison of their centroids. The result of PCC between a pair of cluster centroids of dimension $|\mathbf{O}|$, where \mathbf{O} is the set of observed variables, can be represented by a binary vector of size $|\mathbf{O}|$ in which each element is 1 or 0 depending, respectively, on whether or not there is a difference between the corresponding elements in the compared centroids.

When PCC is between clusters that represent observed major value configurations (i.e., PCC between major clusters), an element of 1 identifies an observed variable that has changed its value between the

compared clusters due to a change in \mathbf{ex} . Thus, the 1s in a major-major PCC provide evidence of causal relationships between \mathbf{EX} and \mathbf{O} . Practically, LPCC always identifies all observed variables that are represented by 1s together in all PCCs as the observed descendants of the same exogenous variable (Section 4.1). However, due to the probabilistic nature of BN and the existence of endogenous latents (mediating the connections from \mathbf{EX} to \mathbf{O}), some of the clusters are k -order minor clusters (in different orders), representing k -order minor configurations/effects. Minor clusters are more difficult to identify than major clusters because the latter reflect the major effects of \mathbf{EX} on \mathbf{EN} and, therefore, are considerably more populated by data points than the former. Nevertheless, minor clusters are important in casual discovery by LPCC even though a major-minor PCC cannot tell the effect of \mathbf{EX} on \mathbf{EN} because an observed variable in two compared (major and minor) clusters should not necessarily change its value as a result of a change in \mathbf{ex} . Their importance is because a major cluster, which is a zero-order minor value configuration and thus has zero minor values, cannot indicate (when compared with another major cluster) the existence of minor values. On the contrary, PCC between major and minor clusters shows (through the number of 1s) the number of minor values represented in the minor cluster, and this is exploited by LPCC for identifying the endogenous latents and interrelations among them (Section 4.4). That is, PCC is the source to identify causal relationships in the unknown LVM; major-major PCCs are used for identifying the exogenous variables and their descendants, and major-minor PCC are used for identifying the endogenous latents, their interrelations, and their observed children.

4. OVERVIEW OF LPCC

To demonstrate the relations between clustering results and learning an LVM, we use G1 in Figure 1 that has two exogenous variables, L1 and L2, each having three observed children [assume all variables are binary, i.e., L1 and L2 have four possible \mathbf{exs} (L1L2= 00, 01, 10, 11)]. We generated a synthetic data set of 1,000 patterns from G1 over the six observed variables, used a uniform distribution over L1 and L2, and set the probabilities of an observed child, X_i , $i = 1, \dots, 6$, given its latent parent, L_k , $k = 1, 2$, to be $P(X_i = v | L_k = v) = 0.8, v = 0, 1$. Using the self-organizing map (SOM) (Kohonen, 1997), we clustered the data set and found 16 clusters, of which four were major (see below how to identify major clusters). This meets our expectation of four major clusters corresponding to the four possible \mathbf{exs} (these clusters are presented in Table 1a in the SMF by their centroids, which are the most prevalent patterns in the clusters, and in Table 1b by their PCCs). For example, $PCC1,2$, comparing clusters $C1$ and $C2$, shows that when moving from $C1$ to $C2$, only the values corresponding to variables X1, X2, and X3 have been

changed (i.e., $\delta X_1 = \delta X_2 = \delta X_3 = 1$ in Table 1b). Lemma 2 guarantees that the three variables are descendants of the same EX that changed its value between two \mathbf{exs} represented by $C1$ and $C2$. $PCC1,4$, $PCC2,3$, and $PCC3,4$ enforce this conclusion. Indeed, we know from the true graph, G1, that this EX is latent L1. A similar conclusion can be deduced about X4, X5, and X6, as descendants of an exogenous latent, which we know, is L2.

LPCC is fed by data that is sampled from the observed variables in the unknown model (Figure 2 in the SMF). It clusters the data using SOM (any other clustering algorithm is good as well), and selects an initial set of major clusters. Then, LPCC learns LVM in two stages. In the first stage, LPCC identifies exogenous latent variables and latent colliders (without distinguishing them yet) and their corresponding observed descendants before distinguishing them. LPCC iteratively improves the selection of the major clusters, and the entire stage is repeated until convergence. In the second stage, LPCC identifies endogenous latent non-colliders with their children. Because this stage cannot distinguish from the outset between latent non-colliders and their latent ancestors, LPCC also needs to apply a mechanism to split these two types of latent variables from each other and to find the links between them after the split.

4.1 IDENTIFICATION OF EXOGENOUS LATENT VARIABLES AND LATENT COLLIDERS

Table 1b shows that $PCC1,2$ (and $PCC3,4$) provides evidence that X1, X2, and X3 may be descendants of the same exogenous latent (L1, as we know) that has changed its value between the two \mathbf{exs} represented by $C1$ and $C2$. Relying only on one PCC may be inadequate when concluding that these variables are descendants of the same exogenous latent because there may be other exogenous latents that have changed their values too. Table 1b shows that $PCC2,3$ (and $PCC1,4$) provides the same evidence about X1, X2, and X3. But, $PCC2,3$ and $PCC1,4$ also show that the values corresponding to X4, X5, and X6 have been changed together too, whereas these values did not change in $PCC1,2$ and $PCC3,4$. Because X4, X5, and X6 changed their values only in $PCC2,3$ but not in $PCC1,2$, they cannot be descendants of L1. This insight strengthens the evidence that X1, X2, and X3 are the only descendants of L1. A similar analysis using $PCC1,3$ and $PCC2,4$ will identify that X4, X5, and X6 are descendants of another latent variable (L2, as we know). Thus, we define (Definition 16) an **MSO** as the *maximal set of observed variables* that always changes its values together in each major-major PCC in which at least one of the variables changes value. For example, X1 (Table 1) changes its value in $PCC1,2$, $PCC1,4$, $PCC2,3$, and $PCC3,4$ and always together with X2 and X3 (and vice versa). Thus {X1, X2, X3} (and similarly {X4, X5, X6}) is an **MSO**. Each **MSO** includes descendants of the same latent variable L, and after considering all PCCs, LPCC learns **MSOs** for all latent variables.

Based on any identified **MSO**, LPCC introduces to the learned graph a new latent variable L together with all the observed variables that are included in this **MSO** as its children. At this stage, LPCC cannot yet distinguish between exogenous latents and latent colliders since the main goal at this stage is to identify latent variables. For now, LPCC focuses on the identification of the relations between the latents and the observed variables, but not on the identification of the interrelations between the latents. The latter task that is needed for distinguishing the collider latents from the exogenous latents is performed in a further step (Section 4.2). Note, however, that the identification of endogenous non-collider latents needs a different analysis that is based on major-minor PCCs and not on major-major PCCs, and thus it is described separately in Section 4.4.

To formalize this identification step (Theorem 1), we 1) define (Definition 17) that a given binary relation (i.e., between two elements) \sim on a set \mathbf{A} is said to be an *equivalence relation* if and only if it is reflexive ($a \sim a$), symmetric (if $a \sim b$ then $b \sim a$), and transitive (if $a \sim b$ and $b \sim c$, then $a \sim c$) for all a, b , and c in \mathbf{A} . The *equivalence class* of a under \sim , denoted $[a]$, is defined as: $[a] = \{b \in \mathbf{A} | b \sim a\}$ (Enderton, 1977); and 2) prove (Lemma 3 in SMF) that the relation “always changes together with” on the set \mathbf{O} of all observed variables, such as “variable $O_i \in \mathbf{O}$ always changes together with variable $O_j \in \mathbf{O}$ in each PCC in which either O_i or O_j has changed” is an equivalence relation. Each equivalence class for this relation comprises an **MSO**. Theorem 1 guarantees that each of multiple latent variables (an exogenous or any of its non-collider descendants or a collider) is identified by its own **MSO**, regardless of the latent cardinality.

Theorem 1 Variables of a particular **MSO** are children of a particular exogenous latent variable EX or its latent non-collider descendant or children of a particular latent collider C .

4.2 DISTINGUISHING LATENT COLLIDERS

After identifying the exogenous latents and latent colliders together, we need to separate them. To demonstrate that, we use graph G_2 in Figure 1, which shows two exogenous latent variables, L_1 and L_3 , that collide in one endogenous latent variable, L_2 (assume all latent variables are binary and each has three binary observed children). Having two exogenous variables, we expect to find four major clusters in the data generated from G_2 ; each corresponds to one of the four possible **ex**s ($L_1L_3 = 00, 01, 10, 11$). We expect the values of X_1, X_2 , and X_3 to change together in all the PCCs in which the value of L_1 changes, and the values of X_7, X_8 , and X_9 to change together in all the PCCs in which the value of L_3 changes. However, the values of X_4, X_5 , and X_6 will change together with those of X_1, X_2 , and X_3 in part of the PCCs and together with those of X_7, X_8 , and X_9 in the remaining PCCs, but always together in all of the PCCs. This will be evidence that

X_4, X_5 , and X_6 are descendants of the same latent variable (L_2 , as we know), which is a collider of L_1 and L_3 .

To learn that an already learned latent variable L is a collider for a set of other already learned (exogenous) latent ancestor variables $\mathbf{LA} \subset \mathbf{EX}$, LPCC requires that: (1) The values of the children of L will change with the values of descendants of different latent variables in \mathbf{LA} in different parts of major-major PCCs; and (2) The values of the children of L will not change in any PCC unless the values of descendants of at least one of the variables in \mathbf{LA} change. This insures that L does not change independently of latents in \mathbf{LA} that are L 's ancestors. We formalize this identification step in Theorem 2:

Theorem 2 A latent variable L is a collider of a set of latent ancestors $\mathbf{LA} \subset \mathbf{EX}$ only if: (1) The values of the children of L change in different parts of some major-major PCCs each time with the values of descendants of another latent ancestor in \mathbf{LA} ; and (2) The values of the children of L do not change in any PCC unless the values of descendants of at least one of the variables in \mathbf{LA} change too.

4.3 STRATEGY FOR CHOOSING MAJOR CLUSTERS

In this problem of unsupervised identification of latent variables given only observational data, LPCC has to deal with lack of prior information regarding the distribution of each latent variable. Therefore, in its first iteration, LPCC assumes a uniform distribution over the latents and selects the major clusters based only on cluster size, which is the number of patterns clustered by the cluster. Clusters that are larger than the average cluster size are selected as majors. However, this initial selection may generate false negative errors for skewed distributions, where a rare value of a latent will be represented only by small clusters that could not be chosen as majors. Also, the initial selection may perform a false positive error, e.g., as a result of a very weak influence of L on any of its children X_i , i.e., $P(X_i = v_1 | L = v) \cong P(X_i = v_2 | L = v)$. This may lead to splitting a data cluster that represents a configuration in which $L = v$ into two clusters with almost the same size, and, when enough samples exist in both clusters, accepting both as major clusters instead of only one.

To avoid the possible errors due to skewed data and circumstances that undermine identifiability, LPCC decides on major clusters iteratively. After learning a graph based on the initial selection of major clusters based on their sizes, it becomes possible to learn the cardinalities of the latent variables and consequently find all possible **ex**s (Section 4.1). Then, for each **ex**, we can select the most probable cluster given the data and use it as an update to the major cluster that represents this **ex**. Using an EM-style

procedure (Dempster et al., 1977), the set of major clusters can be updated iteratively and probabilistically and augment LPCC to learn more accurate graphs. This process can be repeated until convergence to a final graph (Figure 2). Since the final graph depends on the initial graph, the iterative approach cannot guarantee finding the optimal model, but only improving the initial graph.

4.4 IDENTIFICATION OF LATENT NON-COLLIDERS

Based on major-major PCCs, all the endogenous latent non-colliders that are descendants of an exogenous variable EX were temporarily combined with EX , and all the observed children of these latent non-colliders were temporarily combined with the direct children of EX (see example for $G3$ in Figure 1 in the SMF). LPCC splits latent non-colliders from their previously learned ancestor together with their observed children using major-minor PCCs.

A major-minor PCC is between a major cluster and a first-order minor cluster (1-MC). A 1-MC is a cluster that corresponds to a 1-order minor value configuration (Definitions 13 and 14), which exists when exactly one endogenous variable in \mathbf{EN} (either latent or observed) has a minor local value (Definition 13) as a response to a value $ex \in \mathbf{ex}$ that $EX \in \mathbf{EX}$ has obtained. By analyzing, for each exogenous EX , PCCs between 1-MCs and the major clusters that identified EX , LPCC reveals the existence of the latent non-colliders that were previously combined with EX (Section 4.1). Following that, LPCC splits these non-colliders from EX . We will show that if only one observed variable changes in such PCCs (e.g., $X9$ in $PCC1,3$ in Table 3 in the SMF; $C1$ is major and $C3$ is 1-MC) as a response to ex , then the minor value in the 1-MC is of an observed descendant of EX . And, if two or more observed variables change in such PCCs (e.g., $X7-X9$ in $PCC1,6$ in Table 4 in the SMF; $C1$ is major and $C6$ is 1-MC) as a response to ex , then the minor value in the 1-MC is due to a minor value of a latent non-collider descendant of EX . Thus, PCCs between 1-MCs and major clusters that show a change in the values of two or more observed variables provide evidence to the existence of an NC that should be split from its exogenous ancestor. In the SMF, we describe how LPCC finds the set of 1-MCs and why and how the analysis of the PCCs between 1-MCs and major clusters is used to identify and split latent non-colliders from their exogenous ancestor.

We are interested in PCCs between 1-MCs and major clusters that show two sets of two or more elements corresponding to the observed variables. Variables in each set have the same value, which is different than that of the other set. Following, we infer that each set is of a different latent than the one that was expected to be

sole. We denote such PCC by 2S-PCC (i.e., PCC of “two sets”) and the corresponding 1-MC by 2S-MC (Definition 18 in the SMF). Thus, to identify a latent non-collider that was combined to an exogenous latent EX , we consider only the 2S-PCCs; these PCCs are the result of comparing all the 2S-MCs among the 1-MCs for EX with the major clusters that revealed EX . To formalize the identification step, we prove Lemma 4.

Lemma 4 A latent non-collider NC is a descendant of an exogenous latent variable EX . 2S-PCC is PCC between a “two-set” first-order minor cluster 2S-MC due to a minor value in NC and a major cluster that identified EX . \mathbf{ex}' and \mathbf{ex}'' are two value configurations of \mathbf{EX} that correspond to the compared clusters by 2S-PCC. When:

- EX does not change values between \mathbf{ex}' and \mathbf{ex}'' , all the elements in 2S-PCC corresponding to the observed descendants of the latent ancestors of NC (including EX) show no change (i.e., are 0), whereas the elements corresponding to the observed descendants of NC show a change (i.e., are 1),
- and when
- EX changes values between \mathbf{ex}' and \mathbf{ex}'' , all the elements in 2S-PCC corresponding to the observed descendants of the latent ancestors of NC (including EX) show a change (i.e., are 1), whereas the elements corresponding to the observed descendants of NC show no change (i.e., are 0).

Similar to the **MSO** concept that was introduced for major-major PCCs to identify exogenous latents, it is necessary to introduce also for 2S-PCCs a *maximal set of observed variables* (**2S-MSO**) that always change their values together in all 2S-PCCs. Thus, we define (Definition 19 in SMF) a **2S-MSO** as the *maximal set of observed variables* that always change their values together in all 2S-PCCs. After computing all 2S-PCCs for EX , LPCC learns **2S-MSOs** for all these latent variables and thereby identifies all possible splits for EX . We prove (Theorem 3, SMF) that variables of a particular **2S-MSO** are children of an exogenous latent variable EX or any of its descendant latent non-colliders NC .

After splitting the latent non-collider descendants from their exogenous latent ancestor EX , we identify the links between these latents by proving (Proposition 10, SMF) that in 2S-PCCs in which only the observed children of a single latent change, the latent is 1) EX or its leaf latent non-collider descendant, if the connection is serial; or 2) EX 's leaf latent non-collider descendant, if the connection is diverging.

Lemma 4 guarantees that the observed children of a latent non-collider $NC1$, which is a child of another non-collider $NC2$ (both are descendants of EX), will change in all 2S-PCCs with the observed children of

$NC2$ except in a single additional 2S-PCC due to a minor value of $NC1$.

Theorem 4 A latent non-collider $NC1$ is a direct child of another latent non-collider $NC2$ (both on the same path emerging in EX) only if:

- In all 2S-PCCs for which EX does not change, the observed children of $NC1$ always change with those of $NC2$ and also in a single 2S-PCC without the children of $NC2$; and
- In all 2S-PCCs for which a latent non-collider leaf descendant of EX does not change, the observed children of $NC2$ always change with those of $NC1$ and also in a single 2S-PCC without the children of $NC1$.

LPCC uses Theorem 4 to identify the links between the split latents. In the serial connection, there are only two latents with observed children that change alone in some 2S-PCCs; that is EX and its leaf latent non-collider descendant. However, LPCC cannot distinguish between them and thus finds all the links between these two latents as undirected. In the diverging connection, the observed children of EX never change alone (Proposition 10); thus, every latent with children that change alone in some 2S-PCCs can only be a leaf. Thereby, LPCC can identify the directed links among the latents repeatedly on each of the paths from EX to each of the leaves (Theorem 4). Still, LPCC needs to distinguish between the serial and diverging connections. In the case where the observed children of three or more latents change alone in some 2S-PCC, it is clear that it is a diverging connection. Then, LPCC treats these latents as leaves and returns directed paths from EX to each such leaf. However, in the case in which LPCC identifies that the observed children of exactly two latents change alone in some 2S-PCCs, it applies the analysis proposed in Theorem 4 to each of the latents, and if it obtains the same path with opposite

directions, then LPCC considers it as a serial connection and returns the undirected path; otherwise, it considers it as a diverging connection and returns the two directed paths from EX .

5. SUMMARY AND DISCUSSION

We introduced the LPCC concept and algorithm for learning LVMs:

- 1) LPCC learns the model in an unsupervised learning manner. That is, no assumptions about the number of latent variables and their interrelations and which observed variables are the children of which latents are made;
- 2) LPCC combines learning graphical models with data clustering by using the PCC concept to analyze clustering results of discrete variables for learning LVMs;
- 3) LPCC learns MIM, which is a large subclass of SEM;
- 4) LPCC is not limited to latent-tree models and does not make special assumptions, such as linearity, about the distribution;
- 5) LPCC assumes that the measurement model of the true graph is pure and that each latent has at least two indicators. However, if the true graph is not pure, LPCC learns a pure sub-model of the true model, if one exists;
- 6) LPCC is a two-stage algorithm. First, LPCC learns the exogenous latents and the latent colliders, as well as their observed descendants, by utilizing pairwise comparisons between data clusters in the measurement space that may explain latent causes. Second, LPCC learns the endogenous latent non-colliders and their children by splitting these latents from their previously learned latent ancestors; and
- 7) LPCC learns an equivalence class of the structural model of the true graph.




Figure 1: MIMs that are based on a pure measurement model and structural models of different complexity

References

- D. J. Bartholomew, F. Steele, I. Moustaki, and J. I. Galbraith. *The Analysis and Interpretation of Multivariate Data for Social Scientists* (Texts in Statistical Science Series). Chapman & Hall/CRC Press, Boca Raton, Florida, USA, 2002.
- A. Dempster, N. Laird, and D. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of Royal Statistical Society*, B 39:1–39, 1977.
- G. Elidan, N. Lotner, N. Friedman, and D. Koller. Discovering hidden variables: A structure-based approach. In *Advances in Neural Information Processing Systems* 13:479–485, 2000.
- G. Elidan and N. Friedman. Learning the dimensionality of hidden variables. In *Proceedings of the 17th Conference on Uncertainty in Artificial Intelligence*, pages 144–151, Seattle, Washington, 2001.
- H. B. Enderton. *Elements of Set Theory*. Academic Press, New York, New York, 1977.
- N. Friedman. The Bayesian structural EM algorithm. In *Proceedings of the 14th Conference on Uncertainty in Artificial Intelligence*, pages 129–138, San Francisco, CA, 1998.
- S. Harmeling and C. K. I. Williams. Greedy learning of binary latent trees. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 33:1087–1097, 2011.
- T. Kohonen. *Self-Organizing Maps*. Springer-Verlag Press, New York, New York, 1997.
- J. Pearl. *Causality: Models, Reasoning, & Inference*. Cambridge University Press, New York, New York, 2000.
- J. Pearl and T. Verma. A theory of inferred causation. In *Proceedings of the 2nd International Conference on Principles of Knowledge Representation and Reasoning*, pages 441–452, Cambridge, MA, 1991.
- R. Scheines, P. Spirtes, C. Glymour, C. Meek, and T. Richardson. The TETRAD project: Constraint based aids to causal model specification. Technical Report, Department of Philosophy, Carnegie-Mellon University, Pittsburgh, Pennsylvania, 1995.
- S. Shimizu, T. Inazumi, Y. Sogawa, A. Hyvarinen, Y. Kawahara, T. Washio, P. Hoyer, and K. Bolle. DirectLiNGAM: A direct method for learning a linear non-Gaussian structural equation model. *Journal of Machine Learning Research*, 12:1225–1248, 2011.
- R. Silva, R. Scheines, C. Glymour, and P. Spirtes. Learning the structure of linear latent variable models. *Journal of Machine Learning Research*, 7:191–246, 2006.
- P. Spirtes, C. Glymour, and R. Scheines. *Causation, Prediction, and Search*, 2nd edition. MIT Press, New York, New York, 2000.
- P. Spirtes. Calculation of entailed rank constraints in partially non-linear and cyclic models. In *Proceedings of the 29th Conference on Uncertainty in Artificial Intelligence*, pages 606–615, Bellevue, Washington, 2013.
- Y. Wang, N. L. Zhang, and T. Chen. Latent-tree models and approximate inference in Bayesian networks. *Journal of Artificial Intelligence Research*, 32:879–900, 2008.
- N. Zhang. Hierarchical latent class models for cluster analysis. *Journal of Machine Learning Research*, 5:697–723, 2004.

Pairwise Cluster Comparison for Learning Latent Variable Models – Supplementary Material

Nuaman Asbeh

Industrial Engineering and Management
Ben-Gurion University of the Negev
Beer-Sheva, 84105, Israel

Boaz Lerner

Industrial Engineering and Management
Ben-Gurion University of the Negev
Beer-Sheva, 84105, Israel

Definition 5 A path between two nodes V_1 and V_n in a graph \mathbf{G} is a sequence of nodes $\{V_1, \dots, V_n\}$, such that V_i and V_{i+1} are adjacent in \mathbf{G} , $1 \leq i < n$, i.e., $\{V_i, V_{i+1}\} \in \mathbf{E}$.

Note that a unique set of edges is associated with each given path. Paths are assumed to be simple by definition; in other words, no node appears in a path more than once, and an empty path consists of a single node.

Definition 6 A collider on a path $\{V_1, \dots, V_n\}$ is a node V_i , $1 < i < n$, such that V_{i-1} and V_{i+1} are parents of V_i .

Definition 7 A directed path T_{V_n} from V_1 to V_n in a graph \mathbf{G} is a path between these two nodes, such that for every pair of consecutive nodes V_i and V_{i+1} , $1 \leq i < n$ on the path, there is an edge from V_i into V_{i+1} in \mathbf{E} . V_1 is the source, and V_n is the sink of the path. A directed path has no colliders.

Lemma 1

- 1) Each latent non-collider NC_t has only one exogenous latent ancestor EX_{NC_t} , and there is only one directed path T_{NC_t} from EX_{NC_t} (source) to NC_t (sink).

Note that we use the notation NC_t , rather than S_t , since the Lemma applies to both exogenous and endogenous latent non-colliders.

- 2) Each latent collider C_j is connected to a set of exogenous latent ancestors \mathbf{EX}_{C_j} via a set of directed paths \mathbf{T}_{C_j} from \mathbf{EX}_{C_j} (sources) to C_j (sink).

Proof:

- 1) If the latent non-collider is exogenous, $NC_t = EX_t$, then $EX_{NC_t} = EX_t$, and T_{NC_t} is the empty path consisting of EX_t . For example, $EX_{L3} = L3$ and $T_{L3} = L3$ in Graphs G2 and G5 in Figure 1. If, however, the latent non-collider is endogenous, $NC_t = S_t$, and we assume by contradiction that it has more than one exogenous latent ancestor and thus more than one directed path from each exogenous ancestor to S_t (and according to Assumption 5 none of the paths passes through a collider) that collide at S_t , then S_t is a collider. This is contrary to the assumption that NC_t is a non-collider. That is, EX_{S_t} is the only exogenous latent ancestor of S_t , and T_{S_t} is the only directed path from EX_{S_t} through S_t 's parent Pa_t to S_t . For example, $EX_{L5} = L3$ and $T_{L5} = \{L3, L4, L5\}$ in Graph G5 (Figure 1). Q.E.D.

Note that if S_t has no endogenous latent non-collider ancestors, then $Pa_t = EX_{S_t}$ and T_{S_t} equals the ordered sequence $\{EX_{S_t}, S_t\}$, e.g., $EX_{L4} = L3$ and $T_{L4} = \{L3, L4\}$ in Graph G5 (Figure 1).

- 2) Under Assumption 5, any parent Pa_j of latent collider C_j could be either a latent non-collider or an exogenous latent; in other words, $\mathbf{Pa}_j \subset (\mathbf{NC} \cup \mathbf{EX})$. If Pa_j is a latent non-collider, then it is on the directed path T_{C_j} from EX_{C_j} to C_j ; and if Pa_j is an exogenous latent EX_{C_j} , then it is the source of a directed path T_{C_j} (or more than a single directed path) to C_j . $\mathbf{EX}_{C_j} = \cup EX_{C_j}$ is the set of exogenous ancestors of C_j , and $\mathbf{T}_{C_j} = \cup T_{C_j}$ is the set of directed paths from \mathbf{EX}_{C_j} to C_j . For example, $\mathbf{EX}_{L4} = \{L1, L5\}$ and $\mathbf{T}_{L4} = \{\{L1, L2, L3, L4\}, \{L5, L4\}\}$ in G6 (Figure 1). Q.E.D.

Proposition 1 The joint probability over \mathbf{V} due to value assignment \mathbf{ex} to exogenous set \mathbf{EX} is determined only by this assignment and the BN conditional probabilities.

Proof: Since the underlying model is a BN, the joint probability over \mathbf{V} , which is represented by the BN, is factored according to the local Markov assumption for \mathbf{G} . That is, any variable in \mathbf{V} is independent of its non-descendants in \mathbf{G} conditioned on its parents in \mathbf{G} :

$$P(\mathbf{V}) = \prod_{V_i \in \mathbf{V}} P(V_i | \mathbf{Pa}_i), \quad (1)$$

where \mathbf{Pa}_i are the parents of V_i . It can be factorized under our assumptions as:

$$\begin{aligned} P(\mathbf{V}) &= P(\mathbf{EX}, \mathbf{C}, \mathbf{S}, \mathbf{OEX}, \mathbf{OC}, \mathbf{OS}) = \\ &= \prod_{EX_i \in \mathbf{EX}} P(EX_i) \prod_{C_j \in \mathbf{C}} P(C_j | \mathbf{Pa}_j) \prod_{S_t \in \mathbf{S}} P(S_t | Pa_t) \\ &= \prod_{OEX_m \in \mathbf{OEX}} P(OEX_m | EX_m) \prod_{OC_k \in \mathbf{OC}} P(OC_k | C_k) \prod_{OS_v \in \mathbf{OS}} P(OS_v | S_v), \end{aligned} \quad (2)$$

where \mathbf{Pa}_j are the latent parents of the latent collider C_j , Pa_t is the latent parent of the latent non-collider S_t (in other words, $\mathbf{Pa}_j, Pa_t \subset \mathbf{NC}$), $C_k \in \mathbf{C}$ and $S_v \in \mathbf{S}$ are the latent collider and latent non-collider parents of observed variables OC_k and OS_v , respectively, and $EX_m \in \mathbf{EX}$ is the exogenous latent parent of observed variable OEX_m .

The first product in (2) for assignment \mathbf{ex} is of \mathbf{ex} 's priors. In the other five products, the probabilities are of endogenous latents or observed variables conditioned on their parents, which, based on Lemma 1, are either on the directed paths from \mathbf{EX} to the latents/observed variables or exogenous themselves. Either way, any assignment of endogenous latents or observed variables is a result of the assignment \mathbf{ex} to \mathbf{EX} that is mediated to the endogenous latents/observed variables by the BN probabilities:

$$\begin{aligned} P(\mathbf{V} | \mathbf{EX} = \mathbf{ex}) &= P(\mathbf{EX}, \mathbf{C}, \mathbf{S}, \mathbf{OEX}, \mathbf{OC}, \mathbf{OS} | \mathbf{EX} = \mathbf{ex}) = \\ &= \prod_{EX_i \in \mathbf{EX}} P(EX_i = ex_i) \prod_{C_j \in \mathbf{C}} P(C_j = c_j | \mathbf{Pa}_j = \mathbf{pa}_j^{\mathbf{ex}c_j}) \prod_{S_t \in \mathbf{S}} P(S_t = nc_t | Pa_t = pa_t^{\mathbf{ex}S_t}) \\ &= \prod_{OEX_m \in \mathbf{OEX}} P(OEX_m | EX_m = ex_m) \prod_{OC_k \in \mathbf{OC}} P(OC_k = oc_k | C_k = c_k^{\mathbf{ex}c_k}) \\ &= \prod_{OS_v \in \mathbf{OS}} P(OS_v = os_v | S_v = s_v^{\mathbf{ex}S_v}). \end{aligned} \quad (3)$$

where

- ex_i and ex_m are the values of EX_i and EX_m (the latter is the parent of the m th observed child of the exogenous latents), respectively, in the assignment \mathbf{ex} to \mathbf{EX} ;
- $\mathbf{pa}_j^{\mathbf{ex}c_j}$ is the configuration of C_j 's parents due to configuration $\mathbf{ex}c_j$ of C_j 's exogenous ancestors in \mathbf{ex} ;
- $pa_t^{\mathbf{ex}S_t}$ is the value of S_t 's parent due to the value ex_{S_t} of S_t 's exogenous ancestor in \mathbf{ex} ;
- $c_k^{\mathbf{ex}c_k}$ is the value of OC_k 's collider parent due to the configuration $\mathbf{ex}c_k$ of C_k 's exogenous ancestors in \mathbf{ex} ; and
- $s_v^{\mathbf{ex}S_v}$ is the value of OS_v 's non-collider parent due to the value ex_{S_v} of S_v 's exogenous ancestor in \mathbf{ex} . Q.E.D.

Proposition 2 The major local value $MAV_{EN_i}(\mathbf{pa}'_i)$ of an endogenous variable EN_i given a certain configuration of its parents \mathbf{pa}'_i is also certain.

Proof:

Assumption 6 guarantees that given a certain configuration \mathbf{pa}'_i of \mathbf{Pa}_i , there exists a certain value en'_i of EN_i , such that $P(EN_i = en'_i | \mathbf{Pa}_i = \mathbf{pa}'_i) > P(EN_i = en''_i | \mathbf{Pa}_i = \mathbf{pa}'_i)$ for every other value en''_i of EN_i . From the definition of a major local value, $MAV_{EN_i}(\mathbf{pa}'_i) = en'_i$. Q.E.D.

Based on the proof of Proposition 1, we can quantify the effect of \mathbf{ex} on \mathbf{EN} . For example, a major effect of \mathbf{ex} on \mathbf{EN} can be factorized according to the product of major local effects on \mathbf{EN} (weighted by the product of priors, $P(EX_i = ex_i)$):

$$\begin{aligned}
MAE(\mathbf{ex}) &= \prod_{EX_i \in \mathbf{EX}} P(EX_i = ex_i) \prod_{C_j \in \mathbf{C}} MAE_{C_j}(\mathbf{pa}_j^{\mathbf{ex}C_j}) \prod_{S_t \in \mathbf{S}} MAE_{S_t}(pa_t^{\mathbf{ex}S_t}) \\
&\quad \prod_{OEX_m \in \mathbf{OEX}} MAE_{OEX_m}(ex_m) \prod_{OC_k \in \mathbf{OC}} MAE_{OC_k}(c_k^{\mathbf{ex}C_k}) \prod_{OS_v \in \mathbf{OS}} MAE_{OS_v}(s_v^{\mathbf{ex}S_v}) \\
&= \prod_{EX_i \in \mathbf{EX}} P(EX_i = ex_i) \prod_{C_j \in \mathbf{C}} \max_{c'_j} P(C_j = c'_j | \mathbf{Pa}_j = \mathbf{pa}_j^{\mathbf{ex}C_j}) \\
&\quad \prod_{S_t \in \mathbf{S}} \max_{s'_t} P(S_t = s'_t | Pa_t = pa_t^{\mathbf{ex}S_t}) \prod_{OEX_m \in \mathbf{OEX}} \max_{oex'_m} P(OEX_m = oex'_m | EX_m = ex_m) \\
&\quad \prod_{OC_k \in \mathbf{OC}} \max_{oc'_k} P(OC_k = oc'_k | C_k = c_k^{\mathbf{ex}C_k}) \prod_{OS_v \in \mathbf{OS}} \max_{os'_v} P(OS_v = os'_v | S_v = s_v^{\mathbf{ex}S_v}).
\end{aligned} \tag{4}$$

Definition 10 A *partial effect* on a subset of endogenous variables $\mathbf{EN}' \subseteq \mathbf{EN}$ is the influence of a configuration \mathbf{ex}' of \mathbf{EN}' 's exogenous ancestors $\mathbf{EX}' \subseteq \mathbf{EX}$ on \mathbf{EN}' . We define a *partial major effect* $MAE_{\mathbf{EN}'}(\mathbf{ex}')$ as the largest partial effect of \mathbf{ex}' on \mathbf{EN}' and a *partial minor effect* $MIE_{\mathbf{EN}'}(\mathbf{ex}')$ as any non- $MAE_{\mathbf{EN}'}(\mathbf{ex}')$ partial effect of \mathbf{ex}' on \mathbf{EN}' . A *partial major value configuration* $MAV_{\mathbf{EN}'}(\mathbf{ex}')$ is the \mathbf{en}' of \mathbf{EN}' corresponding to $MAE_{\mathbf{EN}'}(\mathbf{ex}')$; in other words, the most probable \mathbf{en}' due to \mathbf{ex}' , and a *partial minor value configuration* is an \mathbf{en}' corresponding to any $MIE_{\mathbf{EN}'}(\mathbf{ex}')$.

We are interested in representing the influence of exogenous variables on their observed descendants and all the variables in the directed paths connecting them. To do this, we separately analyze the (partial) effect of each exogenous variable on each observed variable for which the exogenous is its ancestor and all the latent variables along the path connecting these two. We distinguish between two cases (both are represented in Lemma 1): (1) Observed descendants in \mathbf{OEX} and \mathbf{OS} that are, respectively, children of exogenous latents and children of latent non-colliders that are linked to their exogenous ancestors, each via a single directed path; and (2) Observed descendants in \mathbf{OC} that are children of latent colliders and linked to their exogenous ancestors via a set of directed paths through their latent collider parents. Thus, we are interested in:

1. The partial effect of a value of exogenous ancestor EX_{NC_v} to non-collider NC_v on any configuration of the set of variables $\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}$, where ONC_v is an observed child of latent non-collider NC_v , and TS_{NC_v} is the set of variables in the directed path T_{NC_v} from EX_{NC_v} to NC_v . The corresponding $MAE_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex_{NC_v})$ and $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex_{NC_v})$ are partial major effect and partial major value configuration, respectively. For example, we may be interested in the partial effect of a value of $EX_{NC_v} = EX_{L5} = L3$ in G5 (Figure 1) on $\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\} = \{TS_{L5} \setminus L3, X13\} = \{L4, L5, X13\}$. Note that we use here the notation NC_v since we are interested in both exogenous and endogenous latent non-colliders. When we are interested in the partial effect on an observed variable in \mathbf{OEX} , its exogenous ancestor (which is also its direct parent) is also the latent non-collider, NC_v , and the effect is not measured on any other variable but this observed variable. This is *Case I*, which is analyzed below;

2. The partial effect of a configuration of exogenous variables \mathbf{EX}_{C_k} to collider C_k on any configuration of the set of variables $\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}$, where OC_k is an observed child of latent collider C_k ¹, and \mathbf{TS}_{C_k} is the set of variables in the set of directed paths \mathbf{T}_{C_k} from \mathbf{EX}_{C_k} to C_k . The corresponding $MAE_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}_{C_k})$ and $MAV_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}_{C_k})$ are partial major effect and partial major value configuration, respectively. For example, we may be interested in the partial effect of a configuration of $\mathbf{EX}_{C_k} = \mathbf{EX}_{L4} = \{L1, L5\}$ in G6 (Figure 1) on $\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\} = \{\{L1, L2, L3, L4\} \setminus \{L1\}, \{L5\} \setminus \{L5\}\}, X11\} = \{L2, L3, L4, X11\}$. This is *Case 2*, which is analyzed below.

Following, we provide detailed descriptions for these partial effects and partial values for observed children of latent non-colliders (*Case 1*) and observed children of latent colliders (*Case 2*) and formalize their properties in Propositions 3–7 to set the stage for Lemma 2.

Case 1: Observed children of latent non-colliders

If the latent non-collider NC_v is exogenous, $NC_v = EX_v$ and $ONC_v = OEX_v$, then, $\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\} = OEX_v$. Thus, the partial effect is simply the local effect, and the partial major effect is the major local effect $MAE_{OEX_v}(ex_v)$. If the latent non-collider NC_v is endogenous, then $NC_v = S_v$ and $ONC_v = OS_v$. Then, all variables in $\{TS_{S_v} \setminus EX_{S_v}, OS_v\}$ are d-separated by EX_{S_v} from $\mathbf{EX} \setminus EX_{S_v}$. For example, $\{L4, L5, X13\}$ in G5 (Figure 1) are d-separated by L3 from L2 and its children. Thus, the effect of \mathbf{ex} on the joint probability distribution (3) can be factored to the: a) joint probability over $\mathbf{EX} = \mathbf{ex}$; b) conditional probabilities of the influenced variables along a specific directed path that ends at OS_v on $EX_{S_v} = ex_{S_v}$ (note that the value ex_{S_t} for all $S_t \in TS_{S_v}$ is the same because $EX_{S_t} = EX_v$ is the same exogenous ancestor of all latent non-colliders on the path to S_v); and c) conditional probabilities of all the remaining variables in the graph on $\mathbf{EX} = \mathbf{ex}$:

$$\begin{aligned} P(\mathbf{V} | \mathbf{EX} = \mathbf{ex}) &= P(\mathbf{EX} = \mathbf{ex}) P(TS_{S_v} \setminus EX_{S_v}, OS_v | EX_{S_v} = ex_{S_v}) \\ &P(\mathbf{V} \setminus \{TS_{S_v} \setminus EX_{S_v}, OS_v\} | \mathbf{EX} = \mathbf{ex}), \end{aligned} \tag{5}$$

in which the second factor corresponds to the partial effect of $EX_{S_v} = ex_{S_v}$ on $TS_{S_v} \setminus EX_{S_v}$ (the latent non-colliders on the path from EX_{S_v} to S_v) and S_v 's observed child, OS_v , and the third factor corresponds to the influence of $\mathbf{EX} = \mathbf{ex}$ on all the other (latent and observed) variables in the graph. We can write the second factor describing the partial effect of the value ex_{S_v} on the values of the variables $TS_{S_v} \setminus EX_{S_v}$ in the directed path from EX_{S_v} to OS_v (including) as:

$$\begin{aligned} &P(TS_{S_v} \setminus EX_{S_v}, OS_v | EX_{S_v} = ex_{S_v}) = \\ &\prod_{S_t \in TS_{S_v} \setminus EX_{S_v}} P(S_t = s_t | pa_t = pa_t^{ex_{S_v}}) P(OS_v = os_v | S_v = s_v^{ex_{S_v}}). \end{aligned} \tag{6}$$

The partial major effect in (4) for this directed path can be written as (note again that $ex_{S_t} = ex_{S_v}$):

$$\begin{aligned} MAE_{\{TS_{S_v} \setminus EX_{S_v}, OS_v\}}(ex_{S_v}) &= MAE_{\{TS_{S_v} \setminus EX_{S_v}\}}(ex_{S_v}) \cdot MAE_{OS_v}(s_v^{ex_{S_v}}) = \prod_{S_t \in TS_{S_v} \setminus EX_{S_v}} MAE_{S_t}(pa_t^{ex_{S_v}}) \cdot \\ &MAE_{OS_v}(s_v^{ex_{S_v}}) \end{aligned} \tag{7}$$

Proposition 3 The $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex_{NC_v})$ corresponding to $MAE_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex_{NC_v})$ is a certain value configuration for each certain value ex_{NC_v} .

¹Throughout the paper, we use a child index for its parent. For example, OC_k 's parent is C_k , although generally, we use the index j for a collider, such as C_j .

Proof:

If the latent non-collider NC_v is exogenous, $NC_v = EX_v$ and $ONC_v = OEX_v$, then $\{TS_{NC_v} \setminus EX_{NC_v}, OEX_v\} = OEX_v$ and the partial major value is the local major value $MAV_{OEX_v}(ex_v)$, which by Proposition 2 is certain for a certain value ex_v .

If the latent non-collider NC_v is endogenous, $NC_v = S_v$ and $ONC_v = OS_v$, then we consider $\{TS_{S_v} \setminus EX_{S_v}, OS_v\}$, which is a set of ordered variables along the directed path T_{S_v} that ends in OS_v . The remainder of the proof is by induction:

Basis: Based on Proposition 2, $MAV_{S_1}(ex_{S_1})$, where S_1 is the first variable in $\{TS_{S_v} \setminus EX_{S_v}, OS_v\}$ and a direct child of EX_{S_v} , given a certain value ex_{S_1} , is also certain.

Step: If the major value of the i th variable, S_i , in the subset $\{TS_{S_v} \setminus EX_{S_v}, OS_v\}$, i.e., $MAV_{S_i}(pa_i^{ex_{S_v}})$, is certain for a certain value $pa_i^{ex_{S_v}}$, then the major value of the $(i+1)$ th variable, S_{i+1} , in the subset (which is S_i 's child), i.e., $MAV_{S_{i+1}}(pa_{i+1}^{ex_{S_v}})$, is by Proposition 2 certain too for a certain value $pa_{i+1}^{ex_{S_v}}$ (which is $MAV_{S_i}(pa_i^{ex_{S_v}})$). Q.E.D.

Note that here we use the notation NC_v rather than S_v since the proposition applies to both exogenous and endogenous latent non-colliders.

Proposition 4 All corresponding values in $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex'_{NC_v})$ and $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex''_{NC_v})$, for two values ex'_{NC_v} and ex''_{NC_v} of EX_{NC_v} , are different.

Proof:

If the latent non-collider NC_v is exogenous, $NC_v = EX_v$ and $ONC_v = OEX_v$, then $\{TS_{NC_v} \setminus EX_{NC_v}, OEX_v\} = OEX_v$, and, by Assumption 7, the corresponding $MAV_{OEX_v}(ex'_v)$ and $MAV_{OEX_v}(ex''_v)$ are different for two values ex'_v and ex''_v .

If the latent non-collider NC_v is endogenous, $NC_v = S_v$ and $ONC_v = OS_v$, then we consider $\{TS_{S_v} \setminus EX_{S_v}, OS_v\}$, which is a set of ordered variables along the directed path T_{S_v} that ends in OS_v . The remainder of the proof is by induction:

Basis: The major local values $MAV_{S_1}(ex'_{S_1})$ and $MAV_{S_1}(ex''_{S_1})$ of the first variable, S_1 , in $\{TS_{S_v} \setminus EX_{S_v}, OS_v\}$ (which is also a direct child of EX_{S_v}) and two values ex'_{S_1} and ex''_{S_1} of EX_{S_v} are different based on Assumption 7.

Step: If the major local values of the i th variable, S_i , in $\{TS_{S_v} \setminus EX_{S_v}, OS_v\}$ and two values ex'_v and ex''_v of EX_{S_v} , i.e., $MAV_{S_i}(ex'_{S_i})$ and $MAV_{S_i}(ex''_{S_i})$, are different, then the major local values of S_{i+1} (S_i 's child) and the two values $MAV_{S_i}(ex'_{S_i})$ and $MAV_{S_i}(ex''_{S_i})$, i.e., $MAV_{S_{i+1}}(pa_{i+1}^{ex'_{S_v}}) = MAV_{S_{i+1}}(MAV_{S_i}(ex'_{S_i}))$ and $MAV_{S_{i+1}}(pa_{i+1}^{ex''_{S_v}}) = MAV_{S_{i+1}}(MAV_{S_i}(ex''_{S_i}))$ are different too based on Assumption 7. Q.E.D.

Here also we use the notation NC_v , since the proposition applies to both exogenous and endogenous latent non-colliders.

So far, we have analyzed the impact of an exogenous variable on a latent non-collider by ‘‘propagating’’ the exogenous (source) impact along the path to the latent non-collider (sink). Propositions 3 and 4 guaranteed, respectively, that a certain value of the exogenous variable is responsible for a certain value of the latent non-collider and different values of the exogenous are echoed through different values of the latent non-collider. Proposition 4 is based on the correspondence between changes in values of a latent non-collider and changes in values of its parent; a correspondence that is guaranteed by Assumption 7 (first part). Propositions 3 and 4 ensure, respectively, the existence and uniqueness of the value a latent non-collider gets under the influence of an exogenous ancestor; one (Proposition 3) and only one (Proposition 4) value of the latent non-collider changes with a change in the value of the exogenous. We formalize this in the following Proposition 5.

Proposition 5 EX_{NC_v} changes values (i.e., has two values ex'_{NC_v} and ex''_{NC_v}) if and only if NC_v changes values in the two corresponding major value configurations: $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex'_{NC_v})$ and $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex''_{NC_v})$.

Proof:

(“if”) Proposition 3 guarantees that NC_v has a certain value in $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex_{NC_v})$ for a certain value ex_{NC_v} of EX_{NC_v} . Thus, if NC_v has different values in two $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex_{NC_v})$, then EX_{NC_v} should also have two corresponding values, say ex'_{NC_v} and ex''_{NC_v} .

(“only if”) Proposition 4 guarantees that NC_v will have different values in $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex'_{NC_v})$ and $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex''_{NC_v})$ for two values ex'_{NC_v} and ex''_{NC_v} of EX_{NC_v} . Thus, if NC_v has only a certain value in two $MAV_{\{TS_{NC_v} \setminus EX_{NC_v}, ONC_v\}}(ex_{NC_v})$, then EX_{NC_v} should have also a certain value in the corresponding two ex_{NC_v} . Q.E.D.

Case 2: Observed children of latent colliders

In the case of an observed variable OC_k that is a child of a latent collider C_k , all variables in $\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}$ are d-separated by \mathbf{EX}_{C_k} from $\mathbf{EX} \setminus \mathbf{EX}_{C_k}$. Thus, the effect of \mathbf{ex} on the joint probability distribution (3) can be factored (similarly to Case 1) to the: a) joint probability over $\mathbf{EX} = \mathbf{ex}$; b) conditional probabilities of the influenced variables along all directed paths that end at OC_k on $\mathbf{EX}_{C_k} = \mathbf{ex}_{C_k}$ (note that all variables along each directed path T_{C_k} are influenced by the same ex_{C_k}); and c) conditional probabilities of all the remaining variables in the graph on $\mathbf{EX} = \mathbf{ex}$:

$$\begin{aligned} P(\mathbf{V} | \mathbf{EX} = \mathbf{ex}) &= P(\mathbf{EX} = \mathbf{ex}) P(\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k | \mathbf{EX}_{C_k} = \mathbf{ex}_{C_k}) \\ &P(\mathbf{V} \setminus \{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\} | \mathbf{EX} = \mathbf{ex}), \end{aligned} \quad (8)$$

in which the second factor corresponds to the partial effect on $\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}$ of \mathbf{EX}_{C_k} , and the third factor corresponds to the partial effect on all variables other than $\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}$. We can decompose the second factor into a product of: a) a product over all directed paths into C_k of a product of partial effects over all variables (excluding C_k) in such a path; b) the partial effect on C_k ; and c) that on its child OC_k :

$$\begin{aligned} P(\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k | \mathbf{EX}_{C_k} = \mathbf{ex}_{C_k}) &= \prod_{TS_{C_k} \in \mathbf{TS}_{C_k}} \{ \prod_{S_t \in TS_{C_k} \setminus \{EX_{C_k}, C_k\}} P(S_t = s_t | Pa_t = pa_t^{ex_{C_k}}) \} \\ &P(C_k = c_k | Pa_k = pa_k^{ex_{C_k}}) P(OC_k = oc_k | C_k = c_k^{ex_{C_k}}). \end{aligned} \quad (9)$$

This factor can be rewritten as:

$$\begin{aligned} P(\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k | \mathbf{EX}_{C_k} = \mathbf{ex}_{C_k}) &= \\ \prod_{TS_{C_k} \in \mathbf{TS}_{C_k}} \{ P(TS_{C_k} \setminus \{EX_{C_k}, C_k\} | EX_{C_k} = ex_{C_k}) \} \\ &P(C_k = c_k | Pa_k = pa_k^{ex_{C_k}}) P(OC_k = oc_k | C_k = c_k^{ex_{C_k}}). \end{aligned} \quad (10)$$

It reflects the partial effects of a configuration \mathbf{ex}_{C_k} on the values of the variables in $\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}\}$ and the values C_k and OC_k get, and thus the partial major effect of the second factor can be represented as:

$$MAE_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}_{C_k}) = \prod_{TS_{C_k} \in \mathbf{TS}_{C_k}} \{ MAE_{TS_{C_k} \setminus \{EX_{C_k}, C_k\}}(ex_{C_k}) \} \cdot MAE_{C_k}(pa_k^{ex_{C_k}}) \cdot MAE_{OC_k}(c_k^{ex_{C_k}}). \quad (11)$$

Proposition 6 The $MAV_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}_{C_k})$ corresponding to $MAE_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}_{C_k})$ is a certain value configuration for each certain value configuration \mathbf{ex}_{C_k} .

Proof:

$\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}$ comprises sets of variables $\{TS_{C_k} \setminus EX_{C_k}, OC_k\}$ along all directed paths through C_k that end at OC_k . We will divide each such set into three subsets $\{TS_{C_k} \setminus \{EX_{C_k}, C_k\}\}$, C_k , and OC_k and consider a value configuration for \mathbf{ex}_{C_k} for each subset separately. First, since no latent collider can be a child of a latent collider (Assumption 5), a value configuration for the subset $\{TS_{C_k} \setminus \{EX_{C_k}, C_k\}\}$ is considered to be identical to a value configuration for $\{TS_{N_{C_k}} \setminus EX_{N_{C_k}}\}$, and thus according to Proposition 3, is a certain value configuration for a certain value ex_{C_k} . Because $MAV_{TS_{C_k} \setminus \{EX_{C_k}, C_k\}}(ex_{C_k})$ is a certain value configuration for a certain ex_{C_k} for each directed path TS_{C_k} that is included in \mathbf{TS}_{C_k} , the product of these value configurations, which corresponds to the product of $MAE_{TS_{C_k} \setminus \{EX_{C_k}, C_k\}}(ex_{C_k})$ in (11), is also certain. Second, since C_k 's parents $\mathbf{Pa}_k \subset \cup_{TS_{C_k} \in \mathbf{TS}_{C_k}} \{TS_{C_k} \setminus \{EX_{C_k}, C_k\}\}$, $\mathbf{pa}_k^{\mathbf{ex}_{C_k}}$ are certain value configurations. Thus, based on Proposition 2, $MAV_{C_k}(\mathbf{pa}_k^{\mathbf{ex}_{C_k}})$ is also a certain value and similarly $MAV_{OC_k}(C_k^{\mathbf{ex}_{C_k}})$ is certain, where $C_k^{\mathbf{ex}_{C_k}} = MAV_{C_k}(\mathbf{pa}_k^{\mathbf{ex}_{C_k}})$. Therefore, all variables in $\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}$ are certain in the major configuration for a certain value configuration \mathbf{ex}_{C_k} . Q.E.D.

We wish to apply the same mechanism as in Case 1 to analyze the impact of more than a single exogenous on a latent collider, but here the impact is propagated toward the collider along more than a single path. To accomplish this, the following Proposition 7 analyzes the effect on a collider of each of its exogenous ancestors by considering the effect of such an exogenous on the corresponding collider's parent (using Proposition 5, similar to Case 1 for a latent non-collider) and then the effect of this parent on the collider itself (using the second part of Assumption 7).

Proposition 7 For every exogenous ancestor $EX_{C_k} \in \mathbf{EX}_{C_k}$ of a latent collider C_k , there are at least two configurations \mathbf{ex}'_{C_k} and \mathbf{ex}''_{C_k} of \mathbf{EX}_{C_k} in which only EX_{C_k} of all \mathbf{EX}_{C_k} changes values when C_k changes values in the two corresponding major value configurations $MAV_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}'_{C_k})$ and $MAV_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}''_{C_k})$.

Proof:

We divide the proof into two parts. In the first part, we prove that for each exogenous ancestor of a latent collider, there are at least two MAVs in which only the collider's parent on the path from the exogenous to the collider (of all collider's parents) changes values together with the exogenous. We are aided in this part of the proof by Proposition 5 after considering the collider's parent as a latent non-collider. In the second part, using Assumption 7, we show that each such collider's parent changes values together with the collider in the same two MAVs in which the parent changes values together with the exogenous. Thereby, we prove that for each exogenous ancestor of a latent collider there are at least two MAVs in which the collider changes values only with this exogenous.

For the first part, Proposition 5 guarantees that any exogenous ancestor EX_{C_k} of a parent $Pa_k \in \mathbf{Pa}_k$ of collider C_k (and thus $EX_{Pa_k} = EX_{C_k}$ and Pa_k is also a latent non-collider) changes its value if and only if Pa_k changes its value in two $MAVs_{\{TS_{Pa_k} \setminus EX_{Pa_k}, OPa_k\}}(ex_{Pa_k})$. By the opposite of Proposition 5, any exogenous ancestor $EX_{C_k}^*$ of a parent $Pa_k^* \in \mathbf{Pa}_k \setminus Pa_k$ of C_k is certain if and only if Pa_k^* is certain in two $MAVs_{\{TS_{Pa_k^*} \setminus EX_{Pa_k^*}, OPa_k^*\}}(ex_{Pa_k^*}^*)$.

For the second part, we know by Assumption 7 (second part) that for every C_k that is a latent collider and for every $Pa_k \in \mathbf{Pa}_k$, there are at least two configurations \mathbf{pa}'_k and \mathbf{pa}''_k of \mathbf{Pa}_k in which only the value of Pa_k is different and $MAV_{C_k}(\mathbf{pa}'_k) \neq MAV_{C_k}(\mathbf{pa}''_k)$. That is, the collider (which is the only variable in MAV_{C_k}) changes values together with each of its parents in at least two parents' configurations.

Combining the two parts, we have proven that a collider changes values following a change in the value of each of its parents in at least two configurations of the parents, when the change of values of this parent is due to a change of values of its exogenous ancestor in two exogenous configurations. This means that the collider changes values with each of its exogenous ancestors in at least two exogenous configurations. That is, for two configurations \mathbf{ex}'_{C_k} and \mathbf{ex}''_{C_k} of \mathbf{EX}_{C_k} in which only EX_{C_k} changes values, there are at least two configurations \mathbf{pa}'_k and \mathbf{pa}''_k of \mathbf{Pa}_k in which $Pa_k \in \mathbf{Pa}_k$ changes values in $MAV_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}'_{C_k})$ and $MAV_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}''_{C_k})$ with EX_{C_k} . Since these values of Pa_k in \mathbf{pa}'_k and \mathbf{pa}''_k also change with values of C_k , C_k changes values with EX_{C_k} in \mathbf{ex}'_{C_k} and \mathbf{ex}''_{C_k} . Therefore, there are at least two configurations \mathbf{ex}'_{C_k} and \mathbf{ex}''_{C_k} of \mathbf{EX}_{C_k} in which only EX_{C_k} has changed values when C_k changes values in the two corresponding major value configurations $MAV_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}'_{C_k})$ and $MAV_{\{\mathbf{TS}_{C_k} \setminus \mathbf{EX}_{C_k}, OC_k\}}(\mathbf{ex}''_{C_k})$. Q.E.D.

Lemma 2

- 1) A latent non-collider NC_v and its observed child ONC_v , both descendants of an exogenous variable EX_{NC_v} , change their values in any two major configurations if and only if EX_{NC_v} has changed its value in the corresponding two configurations of \mathbf{EX} .
- 2) A latent collider C_k and its observed child OC_k , both descendants of a set of exogenous variables \mathbf{EX}_{C_k} , change their values in any two major configurations, only if at least one of the exogenous variables in \mathbf{EX}_{C_k} has changed its value in the corresponding two configurations of \mathbf{EX} .

Proof:

- 1) First (“only if”), by Proposition 3, the major value configuration of a latent non-collider NC_v and its observed child ONC_v , both of which are descendants of an exogenous variable EX_{NC_v} , are certain for any certain ex_{NC_v} . That is, if NC_v and ONC_v changed their values in any two major configurations, it is only because EX_{NC_v} has changed its value in the corresponding two configurations of \mathbf{EX} . Second (“if”), by Proposition 4, the major value configurations of NC_v and ONC_v are changed if EX_{NC_v} has changed its value between two configurations of \mathbf{EX} . Q.E.D.
- 2) By Proposition 6, the major value configuration of a collider C_k and its observed child OC_k , both of which are descendants of a set of exogenous variables \mathbf{EX}_{C_k} , are certain for a certain \mathbf{ex}_{C_k} . That is, if C_k and OC_k changed their values in any two major configurations, it is only because at least one of the variables in \mathbf{EX}_{C_k} also changed its value in the corresponding two configurations of \mathbf{EX} . Q.E.D.

The following two propositions formalize the relationships between the observed major value configurations and the set of possible \mathbf{ex} .

Proposition 8 There is only a single observed major value configuration to each exogenous configuration \mathbf{ex} .

Proof:

Based on Lemma 2, different observed major value configurations can be obtained if and only if there is more than a single exogenous configuration. Thus, an exogenous configuration \mathbf{ex} can only lead to a single observed major value configuration. Q.E.D.

Proposition 9 There are different observed major value configurations to different exogenous configurations \mathbf{ex} s.

Proof:

Assume for the sake of contradiction that two different value configurations \mathbf{ex}_1 and \mathbf{ex}_2 led to the same observed major value configuration. Because the two configurations are different, there is at least one exogenous variable EX' that has different values in \mathbf{ex}_1 and \mathbf{ex}_2 , and based on Assumption 4, EX' has at least two observed children. Based on Assumption 7, each of these children has different values in the two observed major value configurations due to the different value of EX' in \mathbf{ex}_1 and \mathbf{ex}_2 . This is contrary to our assumption that there is only one observed major value configuration. Q.E.D.

Table 1.(a) Centroids of Major Clusters for G1 and (b) PCCs between These Major Clusters

Centroid	$X1$	$X2$	$X3$	$X4$	$X5$	$X6$
$C1$	0	0	0	1	1	1
$C2$	1	1	1	1	1	1
$C3$	0	0	0	0	0	0
$C4$	1	1	1	0	0	0

(a)

PCC	$\delta X1$	$\delta X2$	$\delta X3$	$\delta X4$	$\delta X5$	$\delta X6$
$PCC1,2$	1	1	1	0	0	0
$PCC1,3$	0	0	0	1	1	1
$PCC1,4$	1	1	1	1	1	1
$PCC2,3$	1	1	1	1	1	1
$PCC2,4$	0	0	0	1	1	1
$PCC3,4$	1	1	1	0	0	0

(b)




Figure 2: An overview of the LPCC concept.

Note that every two equivalence classes are either equal or disjoint. Therefore, the set of all equivalence classes of \mathbf{A} forms a partition of \mathbf{A} ; every element of \mathbf{A} belongs to one and only one equivalence class. It follows from the properties of an equivalence relation that: $a \sim b$ if and only if $[a] = [b]$. The following Lemma 3 is important since it shows that each **MSO** is an equivalence class, and thus **MSOs** corresponding to the learned latents are disjoint. At this stage, LPCC learns a set of at least two observed variables corresponding to a specific **MSO** for each latent where none of the observed variables is shared with other **MSOs** for other latents; in other words, a pure measurement model.

Lemma 3 The relation “always changes together with” on the set \mathbf{O} of all observed variables, such as “variable $O_i \in \mathbf{O}$ always changes together with variable $O_j \in \mathbf{O}$ in each PCC in which either O_i or O_j has changed” is an equivalence relation. Each equivalence class for this relation comprises an **MSO**.

Proof:

All three conditions that are required for a binary relation to become equivalence are met:

1. O_i always changes with O_i (trivial).
2. If O_i always changes with O_j , then O_j always changes with O_i .
3. If O_i always changes with O_j , and O_j always changes with O_k , then O_i always changes with O_k .

Thus, the set of observed variables in a model can be represented by a set of equivalence classes for this relation, where each equivalence class includes all the variables that have the same equivalence relation, such as an **MSO**. Q.E.D.

Theorem 1 Variables of a particular **MSO** are children of a particular exogenous latent variable EX or its latent non-collider descendant or children of a particular latent collider C .

Proof:

The proof is divided into two separate cases. In the first case, we show that the children of a particular exogenous latent variable or its non-collider descendant belong to the same **MSO**, and in the second case, we show that the children of a particular collider latent belong to the same **MSO**.

Case 1: MSO of observed children of an exogenous variable or its latent non-collider descendants.

Let \mathbf{ONC}_i^2 (in **OEXUOS**) be a set of observed variables that are children of an exogenous variable EX_i and any of its latent non-collider descendants (if they exist), and let \mathbf{OC}_i be a set of observed variables that are children of latent colliders where each has EX_i as an exogenous ancestor with other exogenous variables. Note that \mathbf{OC}_i may be empty, if EX_i does not have any collider descendants, but \mathbf{ONC}_i is never empty because it includes at least \mathbf{OEX}_i (Assumption 4). Because no observed child can be included in both \mathbf{OC}_i and \mathbf{ONC}_i , these sets are disjoint. Their union, $\mathbf{OV}_i = \mathbf{ONC}_i \cup \mathbf{OC}_i$, includes all the observed variables that are affected by EX_i and thus should change their values when EX_i changes.

- First, by Lemma 2 (first part), any subset of variables in \mathbf{ONC}_i (and thus also \mathbf{ONC}_i itself, which is a maximal set) always changes together in all PCCs that correspond to a change in EX_i and never change together in any other PCC. These variables belong to the same **MSO** that represents EX_i .
- Second, let subset \mathbf{OC}_{ij} of \mathbf{OC}_i contain all variables that (1) have a shared exogenous ancestor EX_j (besides EX_i) and (2) change their values together in at least one PCC, which corresponds to a change only in the value of EX_j . By Lemma 2, the other variables in \mathbf{OV}_i that are not descendants of EX_j do not change in that PCC. Thus, variables in $\mathbf{OC}_{ij} \forall j$ do not belong to the same **MSO** for which variables in \mathbf{ONC}_i belong.

Consequently, variables in \mathbf{ONC}_i will change together only in all PCCs that correspond to a change in EX_i , and therefore, will establish a maximal set of the variables $\mathbf{MSO}_i = \mathbf{ONC}_i$ that corresponds to all and only observed variables that are children of exogenous variable EX_i and its latent non-collider descendants.

² So far, observed variables had their own indices and their parents/ancestors also had these indices. In Theorem 1, the index is associated with the exogenous variable (Case 1) and the collider latent (Case 2), since these are the central subjects of interest here.

Case 2: MSO of observed children of a latent collider.

In this case, it is important to note that different colliders and their children are affected by different sets of exogenous variables. Thus, we assume:

Assumption 8 Latent colliders do not share exactly the same sets of exogenous ancestors.

In case Assumption 8 is violated, for example, if several latent colliders share exactly the same set of exogenous ancestors, LPCC does not identify the latent colliders as separate and learns one collider as the parent of all children of the latent colliders. Nevertheless, we believe this assumption is very realistic.

Let \mathbf{OC}_i be the set of the observed variables that are children of latent collider C_i that is a descendant of a set of exogenous variables \mathbf{EX}_{C_i} . By Lemma 2, any variable in \mathbf{OC}_i should not change in any PCC unless at least one of its exogenous ancestors changes. The sets of variables that should change together with variables in \mathbf{OC}_i if any of the exogenous variables in \mathbf{EX}_{C_i} change is represented by:

$$\mathbf{OV} = \bigcup_{EX_t \in \mathbf{EX}_{C_i}} \mathbf{OV}_t = \bigcup_{EX_t \in \mathbf{EX}_{C_i}} \{\mathbf{ONC}_t \cup \mathbf{OC}_t\} = \bigcup_{EX_t \in \mathbf{EX}_{C_i}} \{\{\mathbf{ONC}_t\} \cup \{\mathbf{OC}_t \setminus \mathbf{OC}_i\}\} \cup \mathbf{OC}_i, \quad (12)$$

where the union is over all exogenous ancestors EX_t of C_i . We separate the proof to include three sets of observed variables: 1) \mathbf{OC}_i , which are children of C_i ; 2) \mathbf{ONC}_t , which are children of an exogenous variable EX_t and any of its latent non-collider descendants; and 3) $\{\mathbf{OC}_t \setminus \mathbf{OC}_i\}$, which are children of latent colliders, other than C_i , that are descendants of EX_t .




Figure 3: LVM with two latent colliders.

For example, for latent collider $C_i = L2$ in Figure 3, $\mathbf{EX}_{L2} = \{L1, L3\}$, $\mathbf{OC}_2 = \{X4, X5, X6\}$, $\mathbf{ONC}_{L1} = \{X1, X2, X3\}$, $\{\mathbf{OC}_{L1} \setminus \mathbf{OC}_2\} = \{\}$, $\mathbf{ONC}_{L3} = \{X7, X8, X9\}$, and $\{\mathbf{OC}_{L3} \setminus \mathbf{OC}_2\} = \{X10, X11, X12\}$.

Following, we analyze the three subsets of \mathbf{OV} , specifically, \mathbf{OC}_i , \mathbf{ONC}_t , and $\{\mathbf{OC}_t \setminus \mathbf{OC}_i\}$, and we show that only variables in \mathbf{OC}_i (or any subset of \mathbf{OC}_i) will always change together, whereas other variables in \mathbf{OV} will not. We analyze the subsets \mathbf{ONC}_t and $\{\mathbf{OC}_t \setminus \mathbf{OC}_i\}$ for each exogenous $EX_t \in \mathbf{EX}_{C_i}$; thus, the analysis is also correct for their union (12).

- \mathbf{OC}_i : By Lemma 2 (second part), any subset of variables in \mathbf{OC}_i always changes together in all PCCs that correspond to a change in at least one exogenous variable in \mathbf{EX}_{C_i} . In addition, none of the variables in \mathbf{OC}_i has an exogenous ancestor that is not in \mathbf{EX}_{C_i} ; therefore, no variable in \mathbf{OC}_i ever changes in any PCC that corresponds to an exogenous variable that is not in \mathbf{EX}_{C_i} . These variables belong to the same **MSO** that represents C_i .
- \mathbf{ONC}_t : We previously showed in Case 1, that each \mathbf{ONC}_t forms an **MSO** that corresponds to a single EX_t , and this is the only exogenous ancestor for \mathbf{ONC}_t . By Lemma 3, an **MSO** is an equivalence class; therefore, no other variable in a subset of \mathbf{OV} (including \mathbf{OC}_i) can be added to \mathbf{ONC}_t and it will still remain an **MSO**. Similarly, no subset of variables in \mathbf{ONC}_t can be added to any subset of \mathbf{OV} to obtain an **MSO**.
- $\{\mathbf{OC}_t \setminus \mathbf{OC}_i\}$: Any subset of $\mathbf{OC}_t \setminus \mathbf{OC}_i$ does not change together with any subset of \mathbf{OC}_i because (Assumption 8) for each variable OC_j in $\mathbf{OC}_t \setminus \mathbf{OC}_i$, there is an exogenous ancestor EX_j that is not an ancestor of variables in \mathbf{OC}_i . Thus, by Proposition 7, OC_j changes its value in a PCC that corresponds to a change only in the value of EX_j , whereas the variables in \mathbf{OC}_i , which are not descendants of EX_j , do not change in that PCC.

Consequently, all and only variables in \mathbf{OC}_i (maximal subset of \mathbf{OC}_i) compose **MSO** $_i$ that changes together in all PCCs that correspond to a change in \mathbf{EX}_{C_i} . Q.E.D.

Theorem 2 A latent variable L is a collider of a set of latent ancestors $\mathbf{LA} \subset \mathbf{EX}$ only if: (1) The values of the children of L change in different parts of some major-major PCCs each time with the values of descendants of another latent ancestor in \mathbf{LA} ; and (2) The values of the children of L do not change in any PCC unless the values of descendants of at least one of the variables in \mathbf{LA} change too.

Proof:

Recall that latent variables that have already been learned are either exogenous or colliders. First, we show that a latent variable L that satisfies (2) has to be a collider for a set of latent ancestors $\mathbf{LA} \subset \mathbf{EX}$ by assuming by contradiction that L is not a collider but an exogenous variable. If L is an exogenous, then there exists at least a single major-major PCC_L that corresponds to two \mathbf{ex} s in which only L changes its value. Thus, in PCC_L , only the values of descendants of L change, whereas descendants of other variables in any sub-set $\mathbf{LA} \subset \mathbf{EX}$ do not change. This is in contrast to (2).

Second, we show that if L satisfies (1), then \mathbf{LA} is the set of L 's exogenous ancestors that collide in L . Let \mathbf{ONC}_i (in \mathbf{OEXUOS}) be the set of observed variables that are children of $LA_i \in \mathbf{LA}$ or children of its latent non-collider descendants. Let \mathbf{OC}_i be the set of children of latent colliders where each has LA_i as its ancestor with other exogenous variables in \mathbf{LA} or not. $\mathbf{OV}_i = \mathbf{ONC}_i \cup \mathbf{OC}_i$ includes all the observed variables that are affected by LA_i and thus may change their values when LA_i changes values. In addition, let \mathbf{OC}_L be the set of children of L . We need to show that if L satisfies (1), then $\mathbf{OC}_L \subset \mathbf{OC}_i$ for each $LA_i \in \mathbf{LA}$. Since LA_i is an ancestor of L , (1) ensures that there exists a PCC in which only the values of descendants of LA_i including \mathbf{OC}_L change, whereas the values of descendants of other variables in $\mathbf{LA} \setminus LA_i$ do not change. Thus, $\mathbf{OC}_L \subset \mathbf{OV}_i$. However, none of the children in \mathbf{OC}_L belongs to \mathbf{ONC}_i , otherwise it would have already been identified (Theorem 1) as a descendant of LA_i . Thus, $\mathbf{OC}_L \subset \mathbf{OC}_i$. Q.E.D.

For an example to demonstrate the need in splitting latent non-colliders from their previously learned ancestor, consider G3 in Figure 1, which shows a serial connection of three latent variables L1, L2, and L3. Assume each of the latents is binary and has three binary observed children. L1 is the only \mathbf{EX} with two possible \mathbf{ex} s (L1= 0, 1), and L2 and L3 are \mathbf{NC} s; L2 is a child of L1 and a parent of L3. We synthetically generated a random data set of 1,000 patterns from G3 over the nine observed variables. We set the probabilities of: 1) L1 uniformly; 2) an observed child $X_i, i = 1, \dots, 9$, given its latent parent $L_k, k = 1, 2, 3$ (only if L_k is a direct parent of X_i , e.g., L1 and X1), as $P(X_i = v | L_k = v) = 0.8, v = 0, 1$; and 3) an endogenous latent $L_j, j = 2, 3$, given its latent parent $L_k, k = 1, 2$ (only if L_k is a direct parent of L_j , e.g., L1 and L2), as $P(L_j = v | L_k = v) = 0.8, v = 0, 1$. Table 2 presents the seventeen largest clusters using their centroids and sizes, from which C1 and C2 were selected as major clusters (initially, C1-C6 were selected, because they are larger than the average cluster size of 21, but then the iterative strategy described in Section 4.3 left only C1 and C2 as major clusters). This meets our expectation of two major clusters corresponding to the two possible \mathbf{ex} s of L1. However, because all the elements in $PCC_{1,2}$ are 1's (compare C1 and C2 in Table 2), the nine observed variables establish a single \mathbf{MSO} and by Theorem 1 are considered descendants of the same exogenous variable. That is, the model G0 learned in the first phase of LPCC has only one exogenous latent variable (i.e., L1), and all of the nine observed descendants are learned as its direct children, which is contrary to G3. Since L2 and L3, which are latent non-colliders that are descendants of L1 in G3, were combined in G0 with L1, LPCC should split them from L1 along with their observed children in order to learn the true graph.

Thus, in the second phase, LPCC tests the assumption that G0 is true. If the assumption is rejected, LPCC infers that an exogenous latent \mathbf{EX} has latent non-collider descendants, which were temporarily joined to \mathbf{EX} in the first phase, and hence splits them from \mathbf{EX} . To be able to reject the assumption about the correctness of G0, and thereby identify a possible split of an exogenous latent \mathbf{EX} , we first define a first-order minor cluster (1-MC).

A 1-MC is a cluster that corresponds to a 1-order minor value configuration (Definitions 13 and 14), which exists when exactly one endogenous variable in \mathbf{EN} (either latent or observed) has a minor local value (Definition 13) as a response to a value $\mathbf{ex} \in \mathbf{ex}$ that $\mathbf{EX} \in \mathbf{EX}$ has obtained. By analyzing, for each exogenous \mathbf{EX} , PCCs between 1-MCs and the major clusters that identified \mathbf{EX} , LPCC reveals the existence of the latent non-colliders that were previously combined with \mathbf{EX} (Section 4.1). Following that, LPCC splits these non-colliders from \mathbf{EX} . We will show that if only one observed variable changes in such PCCs (e.g., X9 in $PCC_{1,3}$ in Table 3; C1 is major and C3 is 1-MC) as a response to \mathbf{ex} , then

Table 2: The Seventeen Largest Clusters for G3 Represented by Their Centroids and Sizes

Centroid	X1	X2	X3	X4	X5	X6	X7	X8	X9	size
C1	1	1	1	1	1	1	1	1	1	49
C2	0	0	0	0	0	0	0	0	0	47
C3	1	1	1	1	1	1	1	1	0	28
C4	0	0	0	0	0	0	0	1	0	24
C5	0	1	0	0	0	0	0	0	0	22
C6	1	1	1	1	1	1	0	0	0	22
C7	0	0	1	0	0	0	0	0	0	21
C8	0	0	0	1	1	1	1	1	1	19
C9	0	0	0	0	0	0	1	1	1	18
C10	1	1	1	0	0	0	0	0	0	16
C11	0	0	0	1	0	0	0	0	0	14
C12	0	0	0	0	0	0	1	0	0	14
C13	1	0	1	1	1	1	1	1	1	14
C14	1	1	1	0	1	1	1	1	1	14
C15	1	0	0	0	0	0	0	0	0	13
C16	1	1	1	1	1	1	0	1	1	12
C17	0	0	0	0	0	1	0	0	0	12

the minor value in the 1-MC is of an observed descendant of EX . And, if two or more observed variables change in such PCCs (e.g., $X7-X9$ in $PCC1,6$ in Table 4; $C1$ is major and $C6$ is 1-MC) as a response to ex , then the minor value in the 1-MC is due to a minor value of a latent non-collider descendant of EX . Thus, PCCs between 1-MCs and major clusters that show a change in the values of two or more observed variables provide evidence to the existence of an NC that should be split from its exogenous ancestor. Following, we describe how LPCC finds the set of 1-MCs. Then, we elaborate why and how the analysis of the PCCs between 1-MCs and major clusters is used to identify and split latent non-colliders from their exogenous ancestor.

To find the set of 1-MCs, LPCC first calculates a threshold on the maximal size of 2-order minor clusters (2-MCs). This threshold represents the maximal size of a minor cluster that corresponds to a 2-order minor value configuration, i.e., a minor cluster that represents exactly two endogenous variables in EN that have minor values (Definition 13). This threshold is an approximation for the maximal probability of having minor values as a response to any ex in exactly two descendants of EX , where all other descendants of EX in EN have major values. This approximation is derived from the product of the maximal minor local effects (Definition B.1 in Appendix B) of two observed descendants of EX and the maximal major local effects (Definition B.1) of the other observed descendants in EN (Appendix B). Thus, the sizes of all 1-MCs lie between the maximal size of a 2-MC (i.e., the threshold) and the minimal size of a major cluster (note that a major cluster is also a zero-order minor cluster corresponding to a zero-order minor value configuration). For example, based on the analysis above, $C2$ is the minimal major cluster in learning G3, and all the fifteen clusters (Table 2) that are smaller than $C2$ and larger than the threshold (calculated as 11), i.e., $C3-C17$, are 1-MCs. Note that this procedure is separately applied to each $EX \in \mathbf{EX}$. That is, for each EX , there is a different set of 1-MCs, each representing a single minor value of a descendant of EX and used to identify this descendant, whereas the other descendants of EX have major values.

Recall that every 1-MC corresponds to a 1-order minor value configuration that is due to exactly a single minor value of either an observed variable O or a latent non-collider NC , where both O and NC are descendants of EX in EN . The main difference between these two cases is that in the former, the minor value in O is reflected only in this value, whereas in the latter, the minor value in NC may affect the values of all descendant latents of NC together with those of all the direct children (observed variables) of NC and its descendant latents. A minor value in O happens according to the smallest probability a value of O can take conditioned on a certain value of O 's direct parent. This happens for each value of the direct parent and does not require a change in EX to happen. From definition, a minor value in O in a 1-order minor value configuration can only happen when all EX 's descendants, except O , obtain major values. Although the mechanism of obtaining a minor value in a latent descendant NC

of EX is similar to that in O , the impact of such a minor value is not locally restricted to NC , as for O , but it simultaneously affects all the descendants (latent and observed) of NC , which again, from definition, obtain major values.

We are only interested in the second case of minor values of NC , because their identification helps split this NC from its ancestor EX to which it was initially combined (Section 4.1). Since the observed variables in both cases are among EX 's descendants, which were already used to identify EX , it is a challenge to distinguish between them. Following, we analyze 1-MCs to identify these two cases and concentrate on the second case.

Case 1: A minor value of an observed variable

When comparing, for a specific EX , two centroids – one of a major cluster and the other of a 1-MC that corresponds to an observed minor value configuration (Definition 11) in which an observed variable O , which is a descendant of EX , has a minor value – we can observe that when:

- EX changes values between two **exs** that correspond to the compared clusters, all observed descendants of EX , except O , change values together,

and when

- EX does not change values between two **exs** that correspond to the compared clusters, the only observed descendant of EX that changes value is O .

Thus, a PCC – between the centroid of such 1-MC and a centroid of any of the major clusters – that shows the same value for all, but one (i.e., O), of the observed descendants of EX (i.e., either 1 if EX changes values in the corresponding **exs** or 0 if it does not) identifies a minor value in O . For example, in Table 3, $PCC1,3$ and $PCC2,3$ of $C3$, which is a 1-MC, with the two major clusters $C1$ and $C2$ (Table 2) show the set of observed variables $X1$ - $X8$ that either change values or not together, whereas the single observed variable $X9$ acts contrariwise. This is evidence that $C3$ is a 1-MC due to exactly a single minor value of an observed variable descendent ($X9$) of $L1$ in $G3$. Such an analysis helps LPCC ignore, on the one hand, observed descendants of $L1$ that cannot reflect minor values in $L1$'s latent (non-collider) descendants, and focus, on the other hand, on the latent descendants that should be split from $L1$, as part of Case 2.

Table 3: PCCs for $C3$ with $C1$ and $C2$ (Table 2) in Learning $G3$

Centroid	$X1$	$X2$	$X3$	$X4$	$X5$	$X6$	$X7$	$X8$	$X9$
$PCC1,3$	0	0	0	0	0	0	0	0	1
$PCC2,3$	1	1	1	1	1	1	1	1	0

Case 2: A minor value of a latent non-collider

The minor value of a latent non-collider NC , which is a descendent of EX , can be reflected only via the values of its observed descendants in an observed minor value configuration that is represented by a certain 1-MC. By definition, all of these observed descendants have major values in this 1-order minor configuration since only NC has a minor value in this configuration. The major value of each of these observed descendants is certain given the minor value of NC (Proposition 2) and different from the certain major value it would have if NC had a major value (Assumption 7) instead of its minor value.

When comparing for a specific EX , two centroids – one of a major cluster and the other of a 1-MC that corresponds to an observed minor value configuration in which a latent non-collider NC , which is a descendant of EX , has a minor value – we can observe that when:

- EX changes values between two **exs** that correspond to the compared clusters, all observed descendants of EX , but observed descendants of NC , change values together,

and when

- EX does not change values between two **exs** that correspond to the compared clusters, the only observed descendants of EX that change values are those of NC .

Thus, a PCC – between the centroid of such 1-MC and a centroid of any of the major clusters – that shows two sets of two or more observed variables, each set having a different value, identifies a minor value in NC . The first set in such a PCC comprises the descendants of NC (with a value of 0 if EX changes values in the corresponding exs or 1 if it does not), and the second set comprises all other observed variables that are descendants of EX , but not NC (with a value of 1 if EX changes values in the corresponding exs or 0 if it does not). For example, $PCC1,6$ and $PCC2,6$ (Table 4) of $C6$, which is a 1-MC, with the two major clusters $C1$ and $C2$ (Table 2), show two sets of observed variables for $G3$. The first set consists of $X1$ – $X6$ and the second of $X7$ – $X9$. This is evidence that $C6$ is a 1-MC due to a minor value of a latent non-collider descendant of $L1$, and $L1$ should be split into two latents (each is responsible for one of the two sets). One latent (which we know is $L3$) is a parent of $X7$, $X8$, and $X9$, and the other latent is a parent of $X1$ – $X6$ (which we will show is also split to $L1$ and $L2$, each with its three children). Distinguishing between Case 1 and Case 2 gives us an instrument to identify latent non-colliders.

Definition 18 2S-PCC is PCC between 1-MC and a major cluster that shows two sets of two or more elements corresponding to the observed variables. Elements in each set have the same value, which is different than that of the other set. Accordingly, this 1-MC is defined as 2S-MC.

Table 4: All 2S-PCCs for $G3$

Centroid	$X1$	$X2$	$X3$	$X4$	$X5$	$X6$	$X7$	$X8$	$X9$
$PCC1,6$	0	0	0	0	0	0	1	1	1
$PCC2,6$	1	1	1	1	1	1	0	0	0
$PCC1,8$	1	1	1	0	0	0	0	0	0
$PCC2,8$	0	0	0	1	1	1	1	1	1
$PCC1,9$	1	1	1	1	1	1	0	0	0
$PCC2,9$	0	0	0	0	0	0	1	1	1
$PCC1,10$	0	0	0	1	1	1	1	1	1
$PCC2,10$	1	1	1	0	0	0	0	0	0

The following Theorem 3 helps formalize this identification step, but to prove this theorem, we first need Lemma 4. Recall that the challenge here is to identify a latent non-collider NC that is a descendant of an exogenous latent EX , but was wrongly combined with this exogenous ancestor. To face this challenge, we need to find a circumstance in which EX and NC are involved that is different than that which led to the inability to distinguish between them. NC could not be distinguished from EX when we analyzed major value configurations. But, although a major value configuration is the most probable configuration (Definition 9), minor value configurations are possible too – according to the probability tables of the latents, each given its direct parent – albeit less likely. A minor value configuration in which only NC takes a minor value (i.e., a first-order minor value configuration) is exactly what we need.³ This is because all NC 's latent ancestors, in the first-order minor value configuration, take the same major values they took in the major value configuration and thus influence their descendants the same. But, the minor value NC takes influences its (latent and observed) descendants differently than the major value NC took in the major value configuration. This influence is revealed in the different values the observed children of NC and its descendants take compared to the values they took when NC had a major value. Since the two value configurations are represented in two corresponding clusters – a major cluster and a 2S-MC for NC – the signature of NC can uniquely be detected by comparing the two clusters using 2S-PCC.⁴

Lemma 4 shows that it is possible to identify NC because: 1) Even when EX leads to major values in all NC 's ancestors (and in most cases also in NC), NC can still take a minor value; and 2) even when EX changes values, leading all NC 's ancestors to change values as well, NC can still keep the same

³ All other first-order minor value configurations (due to other latent variables, which are also EX 's descendants) or k -order minor value configurations (Definition 13) due to EX are irrelevant to the identification of NC , although the former – as will be shown in Theorem 3 – play a role in determining the direct observed children of NC among its observed descendants.

⁴ Any 2S-PCC, which is detected for EX , will point to the NC that corresponds to the 2S-MC that is compared by this 2S-PCC.

(minor) value. Thereby, minor value configurations for NC demonstrate its autonomy, enabling its identification and its split from EX .

Lemma 4 A latent non-collider NC is a descendant of an exogenous latent variable EX . 2S-PCC is PCC between a “two-set” first-order minor cluster 2S-MC due to a minor value in NC and a major cluster that identified EX . \mathbf{ex}' and \mathbf{ex}'' are two value configurations of \mathbf{EX} that correspond to the compared clusters by 2S-PCC. When:

- EX does not change values between \mathbf{ex}' and \mathbf{ex}'' , all the elements in 2S-PCC corresponding to the observed descendants of the latent ancestors of NC (including EX) show no change (i.e., are 0), whereas the elements corresponding to the observed descendants of NC show a change (i.e., are 1),

and when

- EX changes values between \mathbf{ex}' and \mathbf{ex}'' , all the elements in 2S-PCC corresponding to the observed descendants of the latent ancestors of NC (including EX) show a change (i.e., are 1), whereas the elements corresponding to the observed descendants of NC show no change (i.e., are 0).

Proof:

2S-MC represents a 1-order minor configuration of \mathbf{EN} in which only NC has a minor value, and all the other variables in \mathbf{EN} have major values. Thus, when

- EX does not change values between \mathbf{ex}' and \mathbf{ex}'' (i.e., $\mathbf{ex}' = \mathbf{ex}''$), then
 1. the major value configuration of the latent ancestors of NC is the same for both \mathbf{ex} s (Proposition 3), and for each such latent, each of its observed children has the same major local value (Proposition 2) for both \mathbf{ex} s. Thus, all the observed children of the latent ancestors of NC do not change values in both clusters, and all the corresponding elements in 2S-PCC are 0; and
 2. NC may take either a major or minor value in response to $\mathbf{ex}' (= \mathbf{ex}'')$, depending on the probabilities that each of NC 's values takes conditioned on the values NC 's direct parent takes. The result of the first case is a major cluster (NC and both its ancestors and descendants have major values) and that of the second case is 1-MC. Since all NC 's ancestors and descendants have major values, whereas NC has a minor value, this 1-MC is 2S-MC by definition. Using these two clusters, LPCC creates 2S-PCC. Since NC d-separates its descendants (both latents and observed) from its ancestors, the values of NC 's descendants are determined only by NC in a way similar to that which we used to prove Proposition 3. Since we are concerned with the case in which NC takes different values for \mathbf{ex}' and \mathbf{ex}'' , its descendants too have different values in the two corresponding configurations, and following Assumption 7, all of their observed children have different values in the corresponding observed configurations and clusters. Therefore, these children change their values between the clusters, as represented by 1s in the 2S-PCC.
- EX changes values between \mathbf{ex}' and \mathbf{ex}'' , then
 1. by Proposition 4, all the latent ancestors of NC have different values for \mathbf{ex}' and \mathbf{ex}'' , and by Assumption 7, all the observed children of these latents have different values for \mathbf{ex}' and \mathbf{ex}'' . Thus, in any 2S-PCC between two clusters corresponding to \mathbf{ex}' and \mathbf{ex}'' , all the elements that correspond to the observed children of the latent ancestors of NC (including EX) show a change (i.e., are 1); and
 2. NC does not change values between \mathbf{ex}' and \mathbf{ex}'' because if it did, then by Proposition 4, all of its latent descendants have different values for \mathbf{ex}' and \mathbf{ex}'' , and by Assumption 7, all of their observed children have different values in the two corresponding observed configurations. And following, in any 2S-PCC between two clusters corresponding to \mathbf{ex}' and \mathbf{ex}'' , all the elements that correspond to NC and its descendants would show a change (i.e., are 1). But, since as we already showed that all the observed children of the ancestors of NC are equal to 1 in these 2S-PCCs, it is contrary to the definition of a 2S-PCC that needs two sets of two or more elements of different values. Thus, NC cannot change values between \mathbf{ex}' and \mathbf{ex}'' . Following and by Proposition 3, all the latent descendants of NC have certain values for this certain value

of NC in both configurations, and by Proposition 2, all the observed children of these latents have certain values in the corresponding observed configurations. Thus, all the elements in 2S-PCC that correspond to the observed children of NC and its descendants do not show a change (i.e., are 0).

Note that the proof implicitly assumes that NC is on a serial connection emerging from EX . In a diverging connection, all the latent variables that are on the paths other than the one that includes NC can be considered with NC 's ancestors because both the latents on the other paths and NC 's ancestors are d-separated (for these 2S-PCCs) by NC from its descendants. Thus, the analysis proposed above for a serial connection generalizes also to the diverging connection. Q.E.D.

Before moving to Theorem 3, let us illustrate the two cases discussed in Lemma 4 for G3. The “ EX does not change values between \mathbf{ex}' and \mathbf{ex}'' ” case can be demonstrated, for example, when comparing $C1$ and $C6$ (Table 2). In response to $EX(=L1)=1$, NC 's (L3) parent (L2) takes a major value of 1 in both the value configurations of the latent variables in response to $\mathbf{ex}' = \mathbf{ex}''$.⁵ Also, L3 takes a major value of 1 in the configuration that is represented by $C1$, which is one of the two major clusters. But, L3, in response to the same configuration of its latent ancestors (L1 and L2), takes a minor value of 0 in the value configuration that is represented by the 2S-MC $C6$. By comparing $C1$ and $C6$, the corresponding 2S-PCC shows two sets of elements: the first of 0s that correspond to the observed variables $X1$ – $X6$, which do not change values between the clusters, and the second of 1s that correspond to $X7$ – $X9$, which do change values between the clusters. This is the evidence we are looking for that is needed to identify L3.

The “ EX changes values between \mathbf{ex}' and \mathbf{ex}'' ” case can be demonstrated, for example, when comparing $C1$ and $C9$ (Table 2). In response to $EX(=L1)=1$ and $EX(=L1)=0$, NC 's (L3) parent (L2) takes a major value of 1 in response to $L1=1$ and a major value of 0 in response to $L1=0$. In the first instance, L3 takes a major value of 1 to create the major configuration that is represented by $C1$, and in the second instance, L3 takes a minor value of 1 in the value configuration that is represented by the 2S-MC $C9$ (and although the first value is major and second is minor, they are both 1). By comparing $C1$ and $C9$, the corresponding 2S-PCC shows two sets of elements, the first of 1s that correspond to the observed variables $X1$ – $X6$, which changed values between the clusters, and the second of 0s that correspond to $X7$ – $X9$, which did not change values between the clusters. This is additional support to the existence of L3. However, relying only on part of the 2S-PCCs may be inadequate to conclude on all possible splits. For example, $PCC1,8$ and $PCC2,8$ (Table 4) show that $X1$ – $X3$ and $X4$ – $X9$ are children of different latents, but do not suggest the split of $X7$ – $X9$ as $PCC1,6$ and $PCC2,6$ do.

Definition 19 A **2S-MSO** is the *maximal set of observed variables* that always change their values together in all 2S-PCCs.

For example, $X1$ in Table 4 changes its value in $PCC2,6$, $PCC1,8$, $PCC1,9$, and $PCC2,10$ and always together with $X2$ and $X3$ (and the other way around). Thus, $\{X1, X2, X3\}$ and similarly $\{X4, X5, X6\}$ and $\{X7, X8, X9\}$ are **2S-MSOs**. Each **2S-MSO** includes children of the same latent non-collider, which is a descendant of EX , or EX itself. After computing all 2S-PCCs for EX , LPCC learns **2S-MSOs** for all these latent variables and thereby identifies all possible splits for EX . Note that compared to **MSO** (Section 4.1), which is identified in major-major PCCs to reveal exogenous latents, **2S-MSO** is identified in PCCs between 2S-MCs and major clusters to reveal splits of latent non-colliders from the exogenous latent that was previously learned using these major clusters.

Theorem 3 Variables of a particular **2S-MSO** are children of an exogenous latent variable EX or any of its descendant latent non-colliders NC .

Proof:

1. Variables of 2S-MSO that are children of EX

We need, first, to prove that the children of EX always change values together and second, that no other observed child of another latent can always change value with them. First, Lemma 4 guarantees that the observed children of EX always change values together since a value change of EX between the

⁵ Note that the values the three latents take in the two-value configurations can only be inferred from the values their children ($X1$ – $X3$ for L1, $X4$ – $X6$ for L2, and $X7$ – $X9$ for L3) take.

two **exs** corresponds to the compared clusters in all 2S-PCCs of 2S-MCs with the major clusters for *EX*. The remainder of the proof is divided into two cases: 1) a serial connection and 2) a diverging connection. In case 1, there exists at least a single 2S-PCC in which only the observed children of *EX* change. This 2S-PCC is between a major cluster for *EX* and 2S-MC due to a minor value of the direct latent non-collider child NC^6 of *EX* (e.g., $L2$ is the direct latent non-collider child of $L1$ in $G3$).⁷ Thus, only the elements in 2S-PCC that correspond to the observed children of *EX* show a change and are equal to 1 (e.g., $PCC2,10$ in Table 4), which guarantees that the observed children of *EX* establish a **2S-MSO**.

In case 2, the same analysis proposed in case 1 is repeated for each of the direct latent non-collider children of *EX* in each of the paths that emerges from *EX*. Let us use the same notation *NC* for each such direct child in each path in turn. In this case, not only do the observed children of *EX* change each time *EX* changes, but also the observed descendants of the other direct latent non-collider children of *EX* (in all paths except that which includes *NC*) change with *EX*. This shows that the observed children of *EX* change with the observed descendants of the direct latent non-collider children of *EX* (all but the descendants of *NC*), but never together with all of them (as at each time, another *NC* is excluded). This guarantees that the observed children of *EX* establish a **2S-MSO**.

II. Variables of **2S-MSO** that are children of *EX*'s descendant *NC*

In a serial connection, we identify three possible situations in which either *NC*, its latent descendant, or its latent ancestor takes a minor value. In each of these situations, no other latent or observed variable can take a minor value because we focus the analysis on 2S-MC through the evaluation of 2S-PCC between this minor cluster and a major cluster for *EX*. For each of the three situations, *EX* may change its value or not, so we have to consider six cases:

1. 2S-MC is due to a minor value of any of *NC*'s latent non-collider descendants, *NCI*, and *EX* does not change value between two **exs** that correspond to the compared clusters. Then, by Lemma 4 (first part), all of *NCI*'s observed descendants do change values but all the observed children of *NCI*'s latent ancestors, including those of *NC*, do not change values.
2. 2S-MC is due to a minor value of any of *NC*'s latent non-collider descendants, *NCI*, and *EX* changes value between two **exs** that correspond to the compared clusters. Then, by Lemma 4 (second part), all of *NCI*'s observed descendants do not change values but all the observed children of *NCI*'s latent ancestors, including those of *NC*, change values.
3. 2S-MC is due to a minor value of *NC*, and *EX* does not change value between two **exs** that correspond to the compared clusters. Then, by Lemma 4 (first part), all of *NC*'s observed descendants do change values but all the observed children of its ancestors do not.
4. 2S-MC is due to a minor value of *NC*, and *EX* changes value between two **exs** that correspond to the compared clusters. Then, by Lemma 4 (second part), all of *NC*'s observed descendants do not change values but all the observed children of its ancestors do.
5. 2S-MC is due to a minor value of *NC*'s latent non-collider ancestor, *NCI*, and *EX* does not change value between two **exs** that correspond to the compared clusters. Then, by Lemma 4 (first part), all the observed children of *NCI* and of its latent descendants, including those of *NC*, change values.
6. 2S-MC is due to a minor value of *NC*'s latent non-collider ancestor, *NCI*, and *EX* changes

⁶ A) We focus on the latent non-collider *NC* that is the direct child of *EX* since only a minor value this *NC* takes can separate *EX* and its observed children from *NC*'s observed children and the observed children of the remaining latent non-colliders, and partition the elements in the corresponding 2S-PCC into two sets in which the first consists of the observed children of *EX* and the second consists of the observed children of all *EX*'s latent descendants. B) In our circumstances, where at least a single latent non-collider has been combined with *EX*, the existence of such a latent variable is guaranteed. C) It is also guaranteed that the 1-MC due to the minor value of the direct latent child of *EX* is 2S-MC because it cannot be due to an observed variable (see Case 2 above).

⁷ We assume that all possible 1-MCs, including the one corresponding to a minor value of the direct latent non-collider child *NC* of *EX*, are found. Practically, if we err in estimating the threshold on the maximal 2-MC (as described above and in Appendix B), we may miss this 1-MC, but this is an identification issue that does not affect the correctness of the theorem.

value between two **ex**s that correspond to the compared clusters. Then, by Lemma 4 (second part), all the observed children of NCI and of its latent descendants, including those of NC , do not change values.

That is, in all six cases, NC 's observed children change values together; in some 2S-PCCs they change values with observed children of a latent non-collider ancestor of NC and in some other 2S-PCCs with observed children of a latent non-collider descendant of NC . Thus, not only will the set of all the observed children of NC always change values together, but also no observed child of any of NC 's latent non-collider ancestors or descendants can be part of this set. This means that the set of observed children of NC is a maximal set of variables that always change together, i.e., **2S-MSO**.

Note that if NC does not have a latent non-collider descendant or ancestor, then Cases 1 and 2 and Cases 5 and 6, respectively, do not exist. In the special case where NC is a leaf (i.e., does not have a latent descendant), Case 3 guarantees that there exists at least a single 2S-PCC in which only the observed children of NC change.

In a diverging connection, all the latent variables that are on paths other than the one that includes NC can be considered with NC 's ancestors because NC d-separates them all from its descendants. Thus, the same analysis proposed in the serial case also holds in the diverging case. Q.E.D.

After splitting the latent non-collider descendants from their exogenous latent ancestor EX , we need to identify the links between these latents. To identify these links, LPCC exploits the following Proposition 10 and Theorem 4. We will see that in the case of a serial connection, LPCC learns the undirected links among the latents, and in the case of a diverging connection, LPCC learns the directed links among the latents. That is, LPCC learns a pattern over the structural model of \mathbf{G} , which represents a Markov equivalence class of models among the latents. In the special case where \mathbf{G} has no serial connection, LPCC learns the true graph.

Proposition 10 In 2S-PCCs in which only the observed children of a single latent change, the latent is

- EX or its leaf latent non-collider descendant, if the connection is serial; or
- EX 's leaf latent non-collider descendant, if the connection is diverging.

Proof:

We already showed that at least a single 2S-PCC exists in the serial connection case in which only the observed children of EX change (Theorem 3). In addition, in the proof of Theorem 3 (II), we already showed that for any NC that is a latent non-collider descendent of EX , NC 's observed children change values in some 2S-PCCs with observed children of a latent non-collider descendant of NC and in the other 2S-PCCs with observed children of a latent non-collider ancestor of NC , but never alone. A special case in the proof of Theorem 3 is when NC is a leaf. Then, at least a single 2S-PCC exists in which only the children of NC change. Q.E.D.

We will exemplify Proposition 10 using $G3$. Table 4 shows all the 2S-PCCs for $G3$ from which we can identify three **2S-MSOs**: $\{X1, X2, X3\}$, $\{X4, X5, X6\}$, and $\{X7, X8, X9\}$. If we consider only 2S-PCCs due to $C1$ (the first major cluster), $\{X1, X2, X3\}$ change alone in $PCCI, 8$, and $\{X7, X8, X9\}$ change alone in $PCCI, 6$. By Proposition 10, these two **2S-MSOs** are observed children of an exogenous latent variable EX and its leaf latent non-collider descendant. From knowing $G3$, we know that these two latents are $L1$ and $L3$. Note that if more than a single leaf of EX exists (i.e., in the case of a diverging connection emerging from EX), then for each such leaf, there is a 2S-PCC in which only the observed children of this leaf change alone. This will help LPCC to identify a diverging connection and determine EX as the source in all paths leading to the leaves. As a result, LPCC could identify the correct direction of the links among the latents.

Proposition 10 guarantees that we find the source (EX) and sink of a path (or in the diverging connection case only the sinks). To identify the directionality between any two latent non-collider variables on the path between the source and sink, we will need more. To motivate that, suppose that when learning $G3$, we already identified $L1$ as EX and $L3$ as EX 's leaf descendant (Proposition 10), and now we have to split $L2$ from $L1$ using the two major clusters, $C1$ and $C2$ (Table 2), which revealed $L1$, and identify the directionality among these three latent variables. Lemma 4 (first part) guarantees that the observed children of a latent non-collider NCI , which is a child of another non-collider $NC2$ (both are

descendants of EX), will change in all 2S-PCCs with the observed children of $NC2$ except in a single additional 2S-PCC due to a minor value of $NC1$. That is, $NC1$ is identified as a direct child of $NC2$, if the observed children of $NC1$ change in all 2S-PCCs (due to a specific major cluster and when EX does not change value) in which the children of $NC2$ change plus in additional 2S-PCCs in which they change without the children of $NC2$.⁸ In our case, this means that the observed children of $L3$, which is a child of $L2$, will change values in all 2S-PCCs in which the observed children of $L2$ change values, and also in an additional 2S-PCC, which is due to a minor value in $L3$. Indeed, $PCCI,10$ (Table 4), due to $C1$, shows that when EX does not change values and the observed children of $L3$, $\{X7,X8,X9\}$, change values, the observed children of $L2$, $\{X4,X5,X6\}$, also change values. In addition, $PCCI,6$, which is the result of comparing $C1$ and 2-MC $C6$ due to a minor value of $L3$, shows that $\{X7,X8,X9\}$ change values without $\{X4,X5,X6\}$ once. $PCC2,8$ and $PCC2,9$ demonstrate the same, when using major cluster $C2$ instead of $C1$ (and $C9$ is the 2-MC that reveals the minor value of $L3$). This provides an indication that $L3$ is a child of $L2$.

But, Proposition 10 cannot guarantee distinguishing between EX and its leaf latent non-collider descendant (hereby a “leaf”); hence, what if we mistakenly identified them? In the $G3$ example, this means we identified $L3$ as EX and $L1$ as EX ’s leaf. Lemma 4 demonstrates an interplay between EX and NC (and all of its descendants) as presented in 2S-PCCs due to a minor value in NC ; when one of them changes, the other does not and vice versa. Because the leaf is one of NC ’s descendants, Lemma 4 guarantees that the observed children of the leaf do not change if and only if EX changes value. That is, by the second part of Lemma 4, if EX changes, then the observed children of the leaf do not change. Thus, if we find 2S-PCCs that show that the observed children of the leaf do not change, then we have evidence that EX changed. This guarantees that the observed children of a latent non-collider $NC2$ (or EX itself), which is a parent of another non-collider $NC1$, will change in all 2S-PCCs with the observed children of $NC1$, except in a single additional 2S-PCC due to a minor value of $NC2$ (or if $NC2$ is EX). In our case, this means that the observed children of $L1$, which is $L2$ ’s parent, will change values in all 2S-PCCs in which the observed children of $L2$ change values, and also in an additional 2S-PCC. Indeed, $PCCI,9$ (Table 4), due to $C1$, shows that when the leaf does not change value and the observed children of $L1$, $\{X1,X2,X3\}$, change values, the observed children of $L2$, $\{X4,X5,X6\}$, also change values. In addition, $PCCI,8$ shows that $\{X1,X2,X3\}$ change values without $\{X4,X5,X6\}$ once. $PCC2,6$ and $PCC2,10$ demonstrate the same when using major cluster $C2$ instead of $C1$. This provides an indication that $L1$ is a child of $L2$, which is the opposite direction between the two in $G3$. That is, the interplay between EX and its leaf lets LPCC identify the directionality between latent non-colliders on the path between EX and the leaf, and in both directions. This means that LPCC can identify only the undirected links between the latents in the serial case.

In the diverging case, the children of EX never change alone, and every latent that its children change alone in some 2S-PCC is a leaf (Proposition 10). Therefore, by performing an analysis as for the serial case using 2S-PCCs in which the observed children of the leaf do not change for each leaf of the branches of the diverging connection, LPCC can identify the links among the latents in opposite directions on each branch. We formalize this by Theorem 4.

⁸ Note that Lemma 4 makes a clear distinction between NC ’s ancestors (and their observed children) and NC ’s descendants (and their observed children), when NC gets a minor value. That is, all NC ’s ancestors follow EX (and change values or not with it) and all NC ’s descendants follow its change of value. This change of NC “breaks” the influence of EX on the latents on the path emerging from EX and “starts” NC ’s own influence on its latent descendants. And this is what is so important in finding the traces of minor values of endogenous latents through 2S-PCCs, that these traces identify the existence of the latents. Particularly, when EX does not change values and all its descendants get major values, the observed children of $NC1$ and $NC2$ will change together, and it is only a minor value that $NC1$ gets that can make a 2S-PCC in which $NC1$ ’s observed children change without those of $NC2$, and thereby indicate that $NC1$ is $NC2$ ’s child.

Theorem 4 A latent non-collider $NC1$ is a direct child of another latent non-collider $NC2$ (both on the same path emerging in EX) only if:

- In all 2S-PCCs for which EX does not change, the observed children of $NC1$ always change with those of $NC2$ and also in a single 2S-PCC without the children of $NC2$; and
- In all 2S-PCCs for which a latent non-collider leaf descendant of EX does not change, the observed children of $NC2$ always change with those of $NC1$ and also in a single 2S-PCC without the children of $NC1$.

Proof:

- Let $NC1$ and $NC2$ be latent non-collider descendants of EX (both on the same path emerging from EX), and $NC1$ is a direct child of $NC2$. A 2S-PCC may result from a 2S-MC due to a minor value in: 1) a latent ancestor of $NC1$ (including $NC2$ itself), 2) $NC1$, or 3) a latent descendent of $NC1$. In the first type of such 2S-PCC (for which EX does not change), Lemma 4 (first part) guarantees that the children of $NC1$ and $NC2$ change together in (1) and do not change at all in (3), whereas in (2) only the observed children of $NC1$ change. Thus, the children of $NC1$ always change with the children of $NC2$, and in addition also in a single 2S-PCC in which the children of $NC2$ do not change.
- In the second type of such 2S-PCC for which the observed children of the leaf latent non-collider descendant of EX do not change, Lemma 4 (second part) guarantees that EX changes value, and the children of $NC1$ and $NC2$ do not change at all in (1) and change together in (3), whereas in (2) only the observed children of $NC2$ change. Thus, the children of $NC2$ always change with the children of $NC1$, and in addition also in a single 2S-PCC in which the children of $NC2$ do not change. The same analysis is true for both a serial and diverging connection. Q.E.D.