

CHOQUET-DENY GROUPS AND THE INFINITE CONJUGACY CLASS PROPERTY

JOSHUA FRISCH, YAIR HARTMAN, OMER TAMUZ,
AND POOYA VAHIDI FERDOWSI

ABSTRACT. A countable discrete group G is said to be Choquet-Deny if it has a trivial Poisson boundary for every generating probability measure. We show that a finitely generated group G is Choquet-Deny if and only if it is virtually nilpotent. For general countable discrete groups, we show that G is Choquet-Deny if and only if none of its quotients have the infinite conjugacy class property. Moreover, when G is not Choquet-Deny, then there exists a symmetric, finite entropy generating measure that has a non trivial Poisson boundary.

1. INTRODUCTION

Let G be a countable discrete group. A probability measure μ on G is *generating* if its support generates G as a semigroup. A function $f: G \rightarrow \mathbb{R}$ is μ -*harmonic* if $f(k) = \sum_{g \in G} \mu(g)f(kg)$ for all $k \in G$. We say that the *measured group* (G, μ) is *Liouville* if all the bounded μ -harmonic functions are constant; this is equivalent to the triviality of the Poisson boundary $\Pi(G, \mu)$ [15–17] (also called the Furstenberg-Poisson boundary; for formal definitions see also, e.g., Furstenberg and Glasner [14], Bader and Shalom [1], or a survey by Furman [13]).

When G is non-amenable, (G, μ) is not Liouville for every generating μ [17]. Conversely, when G is amenable, then there exists some generating μ such that (G, μ) is Liouville, as has been shown by Kaimanovich and Vershik [26] and Rosenblatt [29]. It is natural to ask for which groups G it holds that (G, μ) is Liouville for *every* generating μ . Such groups are known in the literature as *Choquet-Deny* groups (see, e.g., [18–20]). A related term is that of a *Liouville* group (see, e.g., [4, 10]), which usually is taken to mean that (G, μ) is Liouville for every finitely supported generating μ .

The classical Choquet-Deny Theorem (which was first proved for \mathbb{Z}^d by Blackwell [3]) states that abelian groups are Choquet-Deny [5]; the same holds for virtually nilpotent groups [8]. There are many

examples of amenable groups that are not Choquet-Deny: first examples of countable discrete groups¹ are due to Kaimanovich [24] and Kaimanovich and Vershik [26], including locally finite groups; Erschler shows that finitely generated solvable groups that are not virtually nilpotent are not Choquet-Deny [10], and that even some groups of intermediate growth are not Choquet-Deny [9]. Kaimanovich and Vershik [26, p. 466] conjecture that “Given an exponential group G , there exists a symmetric (nonfinitary, in general) measure with nontrivial boundary.” See Bartholdi and Erschler [2] for additional related results and further references and discussion.

Our main result is a characterization of the countable discrete Choquet-Deny groups. Recall that G has the *infinite conjugacy class* property (ICC) if each of its non-trivial elements has an infinite conjugacy class.

Theorem 1. *A countable discrete group G is Choquet-Deny if and only if it has no ICC quotients. Moreover, when G does have an ICC quotient, then there exists a generating, symmetric, finite entropy probability measure μ on G such that (G, μ) is not Liouville. In particular, if G is finitely generated, then it is Choquet-Deny if and only if it is virtually nilpotent.*

The case of a group with no ICC quotients is straightforward and has been previously observed by Jaworski [21] and Jaworski and Raja [22]. We provide a proof for completeness, in §3. Our contribution is therefore in the proof of the other direction, which appears in §2.

The implication for finitely generated groups is a consequence of the fact that in this class, virtually nilpotent groups are precisely those with no ICC quotients (see [28, Theorem 2] and [7, Theorem 2], and a self-contained proof in [11]). Since exponential groups are not virtually nilpotent, Theorem 1 implies that the above mentioned conjecture of Kaimanovich and Vershik [26] is correct.

A very recent result by three of the authors of this paper shows that a group is strongly amenable if and only if it has no ICC quotients [12]. This implies that a countable discrete G is strongly amenable if and only if (G, μ) is Liouville for every generating μ , paralleling the above mentioned characterization of amenability as equivalent to the existence of such a μ . While the proofs of these two similar results are different, it is natural to ask whether there is some deeper connection between strong amenability and the Choquet-Deny property.

The remainder of the paper consists of §2, in which we prove Proposition 2.1, and §3 in which we prove Proposition 3.1. Each proposition

¹In the Lie group setting, an example of an amenable group that is not Choquet-Deny was already known to Furstenberg [15].

shows one of the directions of Theorem 1, with the former establishing the theorem for groups with ICC quotients, and the latter for groups with no ICC quotients. For the convenience of the reader, we informally explain in §2.1 the main ideas behind the proof of Proposition 2.1.

Acknowledgments. We would like to thank Anna Erschler and Vadim Kaimanovich for many useful comments on the first draft of this paper.

2. GROUPS WITH ICC QUOTIENTS

In this section we prove the following proposition. Recall that a probability measure μ on G is symmetric if $\mu(g) = \mu(g^{-1})$ for all $g \in G$. Its Shannon entropy (or just entropy) is $H(\mu) = -\sum_{g \in G} \mu(g) \log \mu(g)$.

Proposition 2.1. *Let G be a countable discrete ICC group with an ICC quotient. Then there exists a generating, symmetric, finite entropy probability measure μ on G such that $\Pi(G, \mu)$ is non-trivial.*

The main technical effort in the proof of Proposition 2.1 is in the proof of the following proposition.

Proposition 2.2. *Let G be an amenable countable discrete ICC group. There exists a fully supported, symmetric, finite entropy probability measure μ on G and an element of the group $h \in G$, such that*

$$(2.1) \quad \liminf_{m \rightarrow \infty} \|h\mu^{*m} - \mu^{*m}\| > 0.$$

Here μ^{*m} is the m -fold convolution $\mu * \cdots * \mu$. We will prove this Proposition later, and now turn to the proof of Proposition 2.1.

Proof of Proposition 2.1. The case of non-amenable G is known, so assume that G is amenable and has an ICC quotient Q . Applying Proposition 2.2 to Q yields a finite entropy, symmetric measure $\bar{\mu}$ on Q that is fully supported, and satisfies (2.1) for some $h \in Q$.

Since $\bar{\mu}$ has full support and satisfies (2.1), it follows from [25, Theorem 2.1] that $(Q, \bar{\mu})$ has a non-trivial Poisson boundary. Let μ be any symmetric, finite entropy generating probability measure on G that is projected to $\bar{\mu}$; the existence of such a μ is straightforward. Then (G, μ) has a non-trivial Poisson boundary. \square

Before we turn to the proof of Proposition 2.2, we would like to give some of the intuition behind it.

2.1. Ideas behind the proof of Proposition 2.2. In this section we provide a simplified description of the main components of the construction in the proof of Proposition 2.2. Note that some of the statements in this description are imprecise, but still give the correct intuition.

To build our measure μ we fix in a probability measure p on the natural numbers, construct inductively a sequence g_1, g_2, \dots of elements of G , and set $\mu(g_n) = p(n)$. With foresight, we choose $p(n) \sim n^{-5/4}$, which will guarantee some desirable properties, as we explain below.

For simplicity assume that G is finitely generated by $\{a_1, \dots, a_k\}$. To construct our sequence g_1, g_2, \dots , begin by setting $g_1 = a_1, \dots, g_k = a_k$. This guarantees that μ is generating. To ensure that μ^{*m} is distant from $h\mu^{*m}$ for some h and large m , we now explain how we choose h and g_{k+1}, g_{k+2}, \dots so that, after removing a low probability set, we are left with disjoint supports for μ^{*m} and $h\mu^{*m}$.

Assume that we already picked g_1, \dots, g_{n-1} . Choose s_1, \dots, s_m i.i.d. from p , and assume that m is such that, with high probability, all the s_i 's are at most n . Then, a random choice from μ^{*m} will be a product $g_{s_1} \cdots g_{s_m}$ of elements in $\{g_1, \dots, g_n\}$. Our choice of p guarantees that with high probability, if g_n appears in this product, then it appears once. In §2.3 we show that p has the properties that we will require.

Assume first, that when picking m i.i.d. draws from μ , there will be a unique element which is g_n and all the rest will be from $\{g_1, \dots, g_{n-1}\}$. Namely, two typical draws from μ^{*m} are of the form $g_{s_1} \cdots g_n \cdots g_{s_{m-1}}$ and $g_{t_1} \cdots g_n \cdots g_{t_m}$ where all the $s_i, t_i < n$.

For the supports of μ^{*m} and $h\mu^{*m}$ to be disjoint (after removing a small probability set), we want to ensure that

$$h \underbrace{g_{s_1} \cdots g_n}_{b_1} \cdots \underbrace{g_n \cdots g_{s_{m-1}}}_{b_2} \neq \underbrace{g_{t_1} \cdots g_n}_{c_1} \cdots \underbrace{g_n \cdots g_{t_m}}_{c_2}$$

which is equivalent to

$$hb_1g_nb_2 \neq c_1g_nc_2.$$

After moving elements around we need to have $g_n^{-1}c_1^{-1}hb_1g_n \neq c_2b_2^{-1}$. Let $C_n = \{g_1, g_1^{-1}, \dots, g_n, g_n^{-1}, e, h\}$ and $X = (C_{n-1})^{2m+1}$. Note that $c_1^{-1}hb_1 \in X$, and $c_2b_2^{-1} \in X$. We want g_n to satisfy that $g_n^{-1}Xg_n \cap X = \emptyset$. This is impossible in general since the identity might belong to X . However, using induction on n , we show that with high probability $c_1^{-1}hb_1 \neq e$, for an appropriate initial choice of h .

We call g_n a *switching element* for X if $g_n^{-1}(X \setminus \{e\})g_n \cap X = \emptyset$, and prove in §2.2 that for any finite set X in an amenable ICC group, there are infinitely many switching elements.

The case that g_n does not appear in both draws but only in one of them requires additionally that g_n is outside some finite set. Since we know that there are infinitely many switching elements, this is always possible. So we will always have a possible choice for g_n .

To ensure that the measure μ is symmetric, we in fact pick g_n or g_n^{-1} with equal probability. This imposes additional constraints on the choice of g_n . Most importantly, we will need to verify not only that $g_n^{-1}(X \setminus \{e\})g_n \cap X = \emptyset$, but that all the four equations $g_n^{\pm 1}(X \setminus \{e\})g_n^{\pm 1} \cap X = \emptyset$ are satisfied. We call such elements g_n *super-switching elements for X* . In §2.2 we prove that like regular switching elements, there are infinitely many super-switching elements in any amenable ICC group. Hence we will be able to find our g_n , avoiding any finite set.

This completes the description of the basic idea behind our construction. In the formal proof we need to address some additional technical challenges. One problem is that for non-finitely generated groups, there is no guarantee that the sequence g_1, g_2, \dots generates the whole group. For that we enumerate all the elements in G by $\{a_1, a_2, \dots\}$ and choose a_n instead of g_n with very small probability.

Now we turn to the formal proof of these ideas.

2.2. Switching Elements. Here we introduce two notions: switching elements and super-switching elements. We will use these notions in the proof of Proposition 2.2.

Definition 2.3. Let G be a countable group and X be a finite symmetric subset of G .

- We call $g \in G$ a *switching element for X* if for all non-identity $x \in X$ we have $g^{-1}xg \notin X$.
- We call $g \in G$ a *super-switching element for X* if for all non-identity $x \in X$ we have $g^{\pm 1}xg^{\pm 1} \notin X$. That is, if $g^{w_1}xg^{w_2} \notin X$ for all non-trivial $x \in X$ and $w_1, w_2 \in \{-1, +1\}$.

Note that since X is symmetric, for $g \in G$ to be super-switching for X , we just need to have that g and g^{-1} are switching for X , and that $gxg \notin X$ for all non-identity $x \in X$.

Proposition 2.4. *Let G be a countable discrete amenable ICC group and let X be a finite symmetric subset of G . The set of super-switching elements for X is infinite.*

Proof of Proposition 2.4. Fix an invariant finitely additive probability measure d on G .² For $A \subseteq G$, we call $d(A)$ the density of A . We will need the fact that infinite index subgroups have zero density.

Let $C_G(x)$ be the centralizer of a non-identity $x \in X$. Then, since X is finite, there is a finite set of cosets of $C_G(x)$ that includes all $g \in G$ such that $g^{-1}xg \in X$. So, non-switching elements for X are in the union of finitely many cosets of subgroups with infinite index, since G is ICC. This means that the set of non-switching elements for X has zero density, and so the set of switching elements for X has density one, as does the set S of elements $g \in G$ such that both g and g^{-1} are switching for X .

Let T be the set of all super-switching elements for X . Let $A \subseteq G$ be the set of involutions $\{g \in G \mid g^2 = e\}$. If $d(A) > 0$, then $d(A \cap S) > 0$. On the other hand, for any $g \in A \cap S$, since g is switching for X and $g^{-1} = g$, g is super-switching for X . Hence $A \cap S \subseteq T$. This shows that if $d(A) > 0$, then $d(T) \geq d(A \cap S) > 0$, and so we are done.

So, we can assume that $d(A) = 0$. For any $x \in X \setminus \{e\}$ and $y \in X$, let $S_{x,y} = \{g \in S \mid gxg = y\}$. Note that

$$T = S \setminus \bigcup_{\substack{x \in X \setminus \{e\} \\ y \in X}} S_{x,y}.$$

It is thus enough to be shown that each $S_{x,y}$ has zero density. So assume for the sake of contradiction that $d(S_{x,y}) > 0$. Fix $g \in S_{x,y}$. We have the following for all $h \in g^{-1}S_{x,y}$.

$$\begin{aligned} gxg = y = ghxgh &\implies (xg) = h(xg)h \\ &\implies (xg)^{-1}h^{-1}(xg) = h \\ &\implies h = (xg)^{-1}h^{-1}(xg) \\ &= (xg)^{-1}[(xg)^{-1}h^{-1}(xg)]^{-1}(xg) \\ &= (xg)^{-2}h(xg)^2 \\ &\implies h \text{ is in the centralizer of } (xg)^2. \end{aligned}$$

So, the centralizer of $(xg)^2$ includes $g^{-1}S_{x,y}$, which has a positive density. So, the centralizer of $(xg)^2$ has finite index. This implies that $(xg)^2 = e$, since G is ICC, and so only the identity can have a finite index centralizer. Hence $xg \in A$ for all $g \in S_{x,y}$. So $xS_{x,y} \subseteq A$. Hence $S_{x,y}$ also has zero density, which is a contradiction. \square

²The argument in this proof can be reproduced using finitary Følner sequence arguments that do not use the axiom of choice.

2.3. A Heavy-Tailed Probability Distribution on \mathbb{N} . Here we state and prove a lemma about the existence of a probability distribution on \mathbb{N} such that infinite i.i.d. samples from this measure have certain properties. We will use this distribution in the proof of Proposition 2.2.

Lemma 2.5. *There exists a probability distribution p on \mathbb{N} with finite Shannon entropy and the following property: for any $\varepsilon > 0$ there exist constants $K_\varepsilon, N_\varepsilon \in \mathbb{N}$ such that for any natural number $m \geq K_\varepsilon$ there exists an $E_\varepsilon^m \subseteq \mathbb{N}^m$ such that:*

- (1) $p^{\times m}(E_\varepsilon^m) \geq 1 - \varepsilon$, where $p^{\times m}$ is the m -fold product measure $p \times \cdots \times p$.
- (2) For any $s = (s_1, \dots, s_m) \in E_\varepsilon^m$, the maximum of $\{s_1, \dots, s_{K_\varepsilon}\}$ is at most N_ε .
- (3) For any $s = (s_1, \dots, s_m) \in E_\varepsilon^m$ and for any $K_\varepsilon \leq k \leq m$, the maximum of $\{s_1, \dots, s_k\}$ is at least k .
- (4) For any $s = (s_1, \dots, s_m) \in E_\varepsilon^m$ and for any $K_\varepsilon \leq k \leq m$, the maximum of $\{s_1, \dots, s_k\}$ appears in (s_1, \dots, s_k) only once.

Proof. Let p be the following probability measure on \mathbb{N} : $p(n) = cn^{-5/4}$, where $1/c = \sum_{n=1}^{\infty} n^{-5/4}$. It is straightforward to see that p has finite entropy.

Let $s = (s_1, s_2, \dots) \in \mathbb{N}^\infty$ have distribution $p^{\times \infty}$; i.e., s is chosen i.i.d. p . Since each s_i has distribution p , for each $n \in \mathbb{N}$ we have:

(2.2)

$$\mathbb{P}[s_i \geq n] = \sum_{m=n}^{\infty} p(m) = c \sum_{m=n}^{\infty} m^{-5/4} \geq c \int_n^{\infty} x^{-5/4} dx = 4cn^{-1/4}.$$

For $k \geq 1$, let

$$\max_s^k := \max\{s_1, \dots, s_k\}.$$

Let A_k be the event that $\max_s^k < k^2$. We have:

$$\begin{aligned} \mathbb{P}[A_k] &= \mathbb{P}[s_i < k^2 \ \forall i \in \{1, \dots, k\}] \\ &= (1 - \mathbb{P}[s_1 < k^2])^k \\ &\leq (1 - 4c(k^2)^{-1/4})^k \\ &\leq e^{-4ck^{1/2}}. \end{aligned}$$

Since the sum of these probabilities is finite, by Borel-Cantelli we get that

$$\mathbb{P}[A_k \text{ infinitely often}] = 0.$$

This means that we have a random index ind'_s which is almost surely finite, and for $A = \{s \mid \max_s^k \geq k^2 \ \forall k \geq \text{ind}'_s\}$ we have $\mathbb{P}[A] = 1$.

For $k \geq 1$, let

$$\text{next}_s^k := \min\{i > k \mid s_i \geq \max_s^k\}.$$

In other words, next_s^k is the first index $i > k$ for which s_i is at least as large as \max_s^k . Let B_k be the event that

$$s_{\text{next}_s^{\text{ind}'_s + k}} = \max_s^{\text{ind}'_s + k}.$$

This is the event that the first entry after $i = \text{ind}'_s + k$ that is at least as large as \max_s^i , is in fact equal to \max_s^i . We would like to show that B_k only happens for finitely many k . To this end, let $C_k^{i,j}$ be the event that $\text{next}_s^{\text{ind}'_s + k} = i$ and $\max_s^{\text{ind}'_s + k} = j$, and note that for any $i, j \in \mathbb{N}$

$$\mathbb{P}[B_k | C_k^{i,j}] = \mathbb{P}[s_i = j | s_i \geq j] \leq \frac{cj^{-5/4}}{4cj^{-1/4}} = \frac{1}{4j}.$$

The equality follows from the independence of s_i from s_1, s_2, \dots, s_{i-1} , and the inequality is a consequence of (2.2). By the definition of ind'_s , we have that $C_k^{i,j}$ can only have positive probability if $j \geq k^2$. Therefore, and since the union $\cup_{i,j} C_k^{i,j}$ has measure one, we have that $\mathbb{P}[B_k] \leq k^{-2}/4$. So, again by Borel-Cantelli, we get

$$\mathbb{P}[B_k \text{ infinitely often}] = 0.$$

This means that we have a random index ind''_s which is almost surely finite, and for $B = \{s \mid s_{\text{next}_s^k} > \max_s^k \quad \forall k \geq \text{ind}'_s + \text{ind}''_s\}$ we have $\mathbb{P}[B] = 1$.

Let $\text{ind}_s = \max_s^{\text{ind}'_s + \text{ind}''_s} + 1$. Note that ind_s is almost surely finite and for $k \geq \text{ind}_s$ we can prove the following properties for s .

- (1) Since $k > \max_s^{\text{ind}'_s + \text{ind}''_s} \geq (\text{ind}'_s + \text{ind}''_s)^2 \geq \text{ind}'_s$, by the definition of ind'_s , we have that $\max_s^k \geq k^2$.
- (2) We know that $\max_s^k \geq k^2 > \max_s^{\text{ind}'_s + \text{ind}''_s}$. So we have indices

$$\text{next}_s^{\text{ind}'_s + \text{ind}''_s} = i_1 < i_2 < \dots < i_l < i_{l+1} = \text{next}_s^k$$

such that $i_j \in \{1, \dots, k\}$ for $j = 1, 2, \dots, l$, $i_{l+1} > k$, and $i_j = \text{next}_s^{i_j - 1}$ for $j = 2, 3, \dots, l + 1$.

Since each $i_j > \text{ind}'_s + \text{ind}''_s$, by the definition of ind''_s , for $j = 1, \dots, l$ we get that $s_{i_j} = \max_s^{i_j} = \max_s^{i_j + 1 - 1}$ appears in $(s_1, \dots, s_{i_j + 1 - 1})$ only once. In particular $\max_s^k = s_{i_l}$ appears in (s_1, \dots, s_k) only once.

Fix $\varepsilon > 0$. Since ind_s is almost surely finite, there are constants $K_\varepsilon \in \mathbb{N}$ and $N_\varepsilon \in \mathbb{N}$ such that the event $E_\varepsilon = \{\text{ind}_s \leq K_\varepsilon \text{ and } \max_s^{K_\varepsilon} \leq N_\varepsilon\}$ has probability at least $1 - \varepsilon$.

Let $m \geq K_\varepsilon$ be a natural number and E_ε^m be the projection of E_ε to the first m coordinates. We have:

- (1) $p^{\times m}(E_\varepsilon^m) \geq \mathbb{P}[E_\varepsilon] \geq 1 - \varepsilon$.
- (2) Any $(s_1, \dots, s_m) \in E_\varepsilon^m$ has an extension $s = (s_1, s_2, \dots) \in E_\varepsilon \subseteq \mathbb{N}^\infty$. Since $s \in E_\varepsilon$, we have $\max_s^{K_\varepsilon} \leq N_\varepsilon$, which means that the maximum of $\{s_1, \dots, s_{K_\varepsilon}\}$ is at most N_ε .
- (3) Any $(s_1, \dots, s_m) \in E_\varepsilon^m$ has an extension $s = (s_1, s_2, \dots) \in E_\varepsilon \subseteq \mathbb{N}^\infty$. Since $s \in E_\varepsilon$, we have $K_\varepsilon \geq \text{ind}_s$. So for $K_\varepsilon \leq k \leq m$ we have $k \geq \text{ind}_s$. The first property we proved for s gives us $\max_s^k \geq k^2 \geq k$, i.e., the maximum of $\{s_1, \dots, s_k\}$ is at least k .
- (4) Any $(s_1, \dots, s_m) \in E_\varepsilon^m$ has an extension $s = (s_1, s_2, \dots) \in E_\varepsilon \subseteq \mathbb{N}^\infty$. Since $s \in E_\varepsilon$, we have $K_\varepsilon \geq \text{ind}_s$. So for $K_\varepsilon \leq k \leq m$ we have $k \geq \text{ind}_s$. The second property we proved for s gives us that the maximum of $\{s_1, \dots, s_k\}$ appears in (s_1, \dots, s_k) only once.

This completes the proof. \square

2.4. Proof of Proposition 2.2. Let $G = \{a_1, a_2, \dots\}$; $\varepsilon > 0$; $p \in P(\mathbb{N})$, $K_\varepsilon, N_\varepsilon \in \mathbb{N}$, and $E_\varepsilon^m \subseteq \mathbb{N}^m$ be the probability measure, the constants, and the events from Lemma 2.5. To simplify notation let $N = N_\varepsilon$ and $K = K_\varepsilon$. Choose g_1, g_2, \dots, g_N arbitrarily in G . For any $n \leq N$ let $A_n = \{g_n, g_n^{-1}, a_n, a_n^{-1}\}$ and $B_n = \cup_{i \leq n} A_i \cup \{e\}$. Choose $h \in G$ such that $h \notin (B_N)^{2N}$, which is a finite set.

For $n = N + 1, N + 2, \dots$ we will define g_n inductively and set $A_n = \{g_n, g_n^{-1}, a_n, a_n^{-1}\}$, $B_n = \cup_{i \leq n} A_i \cup \{e\}$. For any $n \in \mathbb{N}$ let $C_n = B_n \cup \{h^{-1}, h\}$. Note that A_n , B_n , and C_n are finite and symmetric for any $n \in \mathbb{N}$. Now we explain how we define g_{N+1}, g_{N+2}, \dots . Let $n + 1 > N$, and assume that $g_1, \dots, g_n \in G$ are defined. Let $g_{n+1} \in G$ be a super-switching element for $(C_n)^{2n+1}$ which is not in $(C_n)^{8n+1}$. The existence of such super-switching element is guaranteed by Proposition 2.4 and the fact that $(C_n)^{2n+1}$ is a finite symmetric subset of G and $(C_n)^{8n+1}$ is finite.

For $n \in \mathbb{N}$, define a symmetric probability measure μ_n on A_n with

$$\mu_n = \varepsilon 2^{-n} \left(\frac{1}{2} \delta_{a_n} + \frac{1}{2} \delta_{a_n^{-1}} \right) + (1 - \varepsilon 2^{-n}) \left(\frac{1}{2} \delta_{g_n} + \frac{1}{2} \delta_{g_n^{-1}} \right).$$

Here δ_g is the point mass on $g \in G$. Finally, let

$$\mu = \sum_{n=1}^{\infty} p(n) \mu_n.$$

Obviously μ is symmetric and has full support. Since p has finite entropy and each μ_n has support of size at most 4, it follows easily that μ has finite entropy.

We want to show that for μ and h we have

$$\liminf_{m \rightarrow \infty} \|h\mu^{*m} - \mu^{*m}\| > 0.$$

Fix $m \in \mathbb{N}$ larger than K and N . For each $n \in \mathbb{N}$ define $f_n : \{1, 2, 3, 4\} \rightarrow A_n$ by

$$f_n(1) = a_n, f_n(2) = a_n^{-1}, f_n(3) = g_n, f_n(4) = g_n^{-1}.$$

Let

$$\Omega = \{(s, w) \mid s \in \mathbb{N}^m, w \in \{1, 2, 3, 4\}^m\}.$$

Let $\nu : \Omega \rightarrow [0, 1]$ be defined by

$$\nu(s, w) = p^{\times m}(s) \mu_{s_1}(f_{s_1}(w_1)) \mu_{s_2}(f_{s_2}(w_2)) \dots \mu_{s_m}(f_{s_m}(w_m)).$$

Obviously ν is a probability measure on Ω .

Define $r : \Omega \rightarrow G$ by

$$r(s, w) = f_{s_1}(w_1)f_{s_2}(w_2)\dots f_{s_m}(w_m).$$

It is not difficult to see that $r_*\nu = \mu^{*m}$, and so we need to show that $\|hr_*\nu - r_*\nu\|$ is bounded away from zero for large m .

Recall that $E_\varepsilon^m \subseteq \mathbb{N}^m$ is the event given by Lemma 2.5. Fix $s \in E_\varepsilon^m$. Define

$$\begin{aligned} i_{s,1} &= \min\{j \in \{1, \dots, m\} \mid s_j > N\}, \\ i_{s,2} &= \min\{j > i_{s,1} \mid s_j \geq s_{i_{s,1}}\}, \\ &\vdots \\ i_{s,l(s)} &= \min\{j > i_{s,l(s)-1} \mid s_j \geq s_{i_{s,l(s)-1}}\} = \max\{s_1, \dots, s_m\}. \end{aligned}$$

So, $i_{s,1}$ is the index of the first entry of (s_1, \dots, s_m) which is larger than N , $i_{s,2}$ is the index of the next entry that is at least as large, etc. Note that by the second property of E_ε^m in Lemma 2.5, we know that

$$K < i_{s,1} < i_{s,2} < \dots < i_{s,l(s)},$$

and by the fourth property,

$$N < s_{i_{s,1}} < s_{i_{s,2}} < \dots < s_{i_{s,l(s)}}.$$

Let $W_\varepsilon^s = \{w \in \{1, 2, 3, 4\}^m \mid \forall k \leq l(s) \ w_{i_{s,k}} = 3, 4\}$. Note that for $\Omega^s = \{s\} \times \{1, 2, 3, 4\}^m \subseteq \Omega$, we have

$$\begin{aligned} \mathbb{P}[W_\varepsilon^s | \Omega^s] &= 1 - \mathbb{P}[\neg W_\varepsilon^s | \Omega^s] \\ &= 1 - \mathbb{P}\left[w_{i_{s,1}} = 1, 2; \text{ or } w_{i_{s,2}} = 1, 2; \dots; \text{ or } w_{i_{s,l(s)}} = 1, 2 \mid \Omega^s\right] \\ &\geq 1 - \sum_{k=1}^{l(s)} \mathbb{P}[w_{i_{s,k}} = 1, 2 | \Omega^s] \\ &= 1 - \sum_{k=1}^{l(s)} \varepsilon 2^{-s_{i_{s,k}}} \\ &\geq 1 - \sum_{j=1}^{\infty} \varepsilon 2^{-j} = 1 - \varepsilon. \end{aligned}$$

The last inequality is because $s_{i_{s,1}} < s_{i_{s,2}} < \dots < s_{i_{s,l(s)}}$.

Finally, let

$$\Omega_\varepsilon = \{(s, w) \in \Omega \mid s \in E_\varepsilon^m, w \in W_\varepsilon^s\},$$

and note that

$$\nu(\Omega_\varepsilon) \geq (1 - \varepsilon)(1 - \varepsilon) > 1 - 2\varepsilon.$$

Claim 2.6. *For any $\alpha, \beta \in \Omega_\varepsilon$, we have $hr(\alpha) \neq r(\beta)$.*

We prove this claim after we finish the proof of the Proposition.

Let ν_1 be equal to ν conditioned on Ω_ε , and ν_2 be equal to ν conditioned on the complement of Ω_ε . We have $\nu = \nu(\Omega_\varepsilon)\nu_1 + (1 - \nu(\Omega_\varepsilon))\nu_2$, and by the above claim we know $\|hr_*\nu_1 - r_*\nu_1\| = 2$. So

$$\begin{aligned} \|h\mu^{*m} - \mu^{*m}\| &= \|hr_*\nu - r_*\nu\| \\ &= \|\nu(\Omega_\varepsilon)(hr_*\nu_1 - r_*\nu_1) + (1 - \nu(\Omega_\varepsilon))(hr_*\nu_2 - r_*\nu_2)\| \\ &\geq \nu(\Omega_\varepsilon) \|hr_*\nu_1 - r_*\nu_1\| - 2(1 - \nu(\Omega_\varepsilon)) \\ &\geq 2(1 - 2\varepsilon) - 2(2\varepsilon) = 2 - 8\varepsilon, \end{aligned}$$

which is bounded away from zero for ε small enough and all m large enough. This completes the proof of Proposition 2.2.

Proof of Claim 2.6. Let $\alpha = (s, w)$, $\beta = (t, v) \in \Omega_\varepsilon$. Hence $\max\{K, N\} < m$, $s \in E_\varepsilon^m$, $t \in E_\varepsilon^m$, $w \in W_\varepsilon^s$, and $v \in W_\varepsilon^t$. Assume that $hr(\alpha) = r(\beta)$. So, we have

$$hf_{s_1}(w_1) \cdots f_{s_m}(w_m) = ft_1(v_1) \cdots ft_m(v_m).$$

Let $K < i_1 < i_2 < \dots < i_{l(s)}$ and $K < j_1 < j_2 < \dots < j_{l(t)}$ be the indices we defined for s and t in the proof of Proposition 2.2. We

remind the reader that the unique maximum of (s_1, \dots, s_m) is attained at $i_{l(s)}$, with a corresponding statement for (t_1, \dots, t_m) and $j_{l(t)}$. So we have

$$\begin{aligned} & h \overbrace{f_{s_1}(w_1) \cdots f_{s_{i_{l(s)}-1}}(w_{i_{l(s)}-1})}^{b_1} f_{s_{i_{l(s)}}}(w_{i_{l(s)}}) \overbrace{f_{s_{i_{l(s)}+1}}(w_{i_{l(s)}+1}) \cdots f_{s_m}(w_m)}^{b_2} \\ &= \underbrace{f_{t_1}(v_1) \cdots f_{t_{j_{l(t)}-1}}(v_{j_{l(t)}-1})}_{c_1} f_{t_{j_{l(t)}}}(v_{j_{l(t)}}) \underbrace{f_{t_{j_{l(t)}+1}}(v_{j_{l(t)}+1}) \cdots f_{t_m}(v_m)}_{c_2}. \end{aligned}$$

Let $p = s_{i_{l(s)}} = \max\{s_1, \dots, s_m\}$ and $q = t_{j_{l(t)}} = \max\{t_1, \dots, t_m\}$. Since $w \in W_\varepsilon^s$ and $v \in W_\varepsilon^t$, we know $f_{s_{i_{l(s)}}}(w_{i_{l(s)}}) = g_p^{\pm 1}$ and $f_{t_{j_{l(t)}}}(v_{j_{l(t)}}) = g_q^{\pm 1}$, so

$$(2.3) \quad hb_1 g_p^{\pm 1} b_2 = c_1 g_q^{\pm 1} c_2.$$

Since $p = \max\{s_1, \dots, s_m\}$, and since $m \geq K$, we know that $m \leq p$. So $b_1, b_2 \in (B_{p-1})^{p-1} \subseteq (C_{p-1})^{p-1}$. Similarly $c_1, c_2 \in (C_{q-1})^{q-1}$.

Consider the case that $p > q$. Then $c_1, c_2, g_q^{\pm 1} \in (C_q)^q \subseteq (C_{p-1})^{p-1}$. Hence $g_p^{\pm 1} = [b_1^{-1} h^{-1} [c_1 g_q^{\pm 1} c_2 b_2^{-1}]]$ by (2.3), and so

$$g_p \in (C_{p-1})^{4(p-1)} \{h, h^{-1}\} (C_{p-1})^{4(p-1)} \subseteq (C_{p-1})^{8(p-1)+1},$$

which is a contradiction with our choice of g_p , since $p > N$. Similarly, if $p < q$, we get a contradiction. So we can assume that $p = q$.

If $p = q$, then by (2.3) we have

$$hb_1 g_p^{\pm 1} b_2 = c_1 g_p^{\pm 1} c_2,$$

and $c_1, c_2, b_1, b_2 \in (C_{p-1})^{p-1}$. So, for $x = c_1^{-1} h b_1 \in (C_{p-1})^{2(p-1)+1}$ we have $g_p^{\pm 1} x g_p^{\pm 1} = c_2 b_2^{-1} \in (C_{p-1})^{2(p-1)} \subseteq (C_{p-1})^{2(p-1)+1}$. By the fact that g_p is a super-switching element for $(C_{p-1})^{2(p-1)+1}$, we get that x is the identity.

So $hb_1 = c_1$, i.e.

$$h f_{s_1}(w_1) \cdots f_{s_{i_{l(s)}-1}}(w_{i_{l(s)}-1}) = f_{t_1}(v_1) \cdots f_{t_{j_{l(t)}-1}}(v_{j_{l(t)}-1}).$$

By the exact same argument, we can see this leads to a contradiction unless

$$h f_{s_1}(w_1) \cdots f_{s_{i_{l(s)}-1}}(w_{i_{l(s)}-1}) = f_{t_1}(v_1) \cdots f_{t_{j_{l(t)}-1}}(v_{j_{l(t)}-1}).$$

And again, this leads to a contradiction unless

$$h f_{s_1}(w_1) \cdots f_{s_{i_{l(s)}-2}}(w_{i_{l(s)}-2}) = f_{t_1}(v_1) \cdots f_{t_{j_{l(t)}-2}}(v_{j_{l(t)}-2}).$$

Note that if $l(s) \neq l(t)$, at some point in this process we get that either all the s_i 's or all the t_i 's are at most N while the other string has characters strictly greater than N . This leads to a contradiction

similar to the case $p \neq q$, which we explained before. So, by continuing this process, we get a contradiction unless

$$h \overbrace{f_{s_1}(w_1) \cdots f_{s_{i_1-1}}(w_{i_1-1})}^{b'} = \overbrace{f_{t_1}(v_1) \cdots f_{t_{j_1-1}}(v_{j_1-1})}^{c'}.$$

- (1) Recall that $s_1, \dots, s_{i_1-1} \leq N$, which means $\max\{s_1, \dots, s_{i_1-1}\} \leq N$.
- (2) On the other hand $s_{i_1} > N$ and we know that $s_1, \dots, s_K \leq N$. So, $i_1 > K \implies i_1 - 1 \geq K$. So, by the third property of E_ε^m in Lemma 2.5, we get that $\max\{s_1, \dots, s_{i_1-1}\} \geq (i_1 - 1)$.
- (3) Combining the last two results we get $N \geq i_1 - 1$.
- (4) Again, since $s_1, \dots, s_{i_1-1} \leq N$, we get that

$$f_{s_1}(w_1), \dots, f_{s_{i_1-1}}(w_{i_1-1}) \in B_N.$$

So,

$$b' = f_{s_1}(w_1) \cdots f_{s_{i_1-1}}(w_{i_1-1}) \in (B_N)^{i_1-1} \subseteq (B_N)^N.$$

A similar argument shows that $c' = f_{t_1}(v_1) \cdots f_{t_{j_1-1}}(v_{j_1-1}) \in (B_N)^N$. But we know that $hb' = c' \implies h = c'b'^{-1} \in (B_N)^{2N}$, which is in contradiction with our choice of h . □

Remark 2.7. *Note that it is possible, by a straightforward modification of this proof, to allow h to be any non-identity element of G .*

3. GROUPS WITH NO ICC QUOTIENTS

In this section we prove the following.

Proposition 3.1. *Let G be a countable discrete group with no ICC quotients. Then $\Pi(G, \mu)$ is trivial for any generating measure μ on G .*

This claim has appeared previously in the literature [22], and we provide a proof for completeness. The following theorem is in particular well known, and in this form is due to Dynkin and Maljutov [8]. We provide a proof that uses an idea of Margulis [27]. Another proof, using the uniqueness of the stationary measure on the Poisson boundary, appears in [23].

Theorem 3.2. *Let G be a group and let μ be any generating probability measure on G . Then the center of G acts trivially on $\Pi(G, \mu)$.*

Proof. Let Z denote the center of G . We show that the restriction to Z of any bounded μ -harmonic on G is constant.

Endow $[0, 1]^Z$ with the product topology, and let $F \subset [0, 1]^Z$ denote the set of all the restriction to Z of μ -harmonic functions which are

bounded by 0 and 1. Then F is a compact convex space. The *Krein-Milman Theorem* states that F is the closure of the convex hull of its extremal points. Hence it is enough to show that the extremal points in F are constant functions, and hence the bounded μ -harmonic functions are Z -invariant.

Consider the G -action on μ -harmonic functions given by $f^g(x) = f(gx)$, and define the Z -action on F similarly. Let f be an extremal function in F . So $f = \bar{f}|_Z$ is the restriction to Z of some μ -harmonic function \bar{f} . Then for any $z \in Z$ we get

$$f(z) = \bar{f}(z) = \sum_g \mu(g) \bar{f}(zg) = \sum_g \mu(g) \bar{f}(gz) = \sum_g \mu(g) \bar{f}^g(z).$$

Since the equation above holds for any $z \in Z$, we have shown that $f = \sum_g \mu(g) \bar{f}^g|_Z$. Since f is extremal, and since $\bar{f}^g|_Z \in F$, $f = \bar{f}^g|_Z$ for all g in the support of μ . By applying the same argument to convolution powers of μ , and since μ is generating, we have that $f = \bar{f}^g|_Z$ for all $g \in G$, and in particular $f = \bar{f}^z|_Z = f^z$ for all $z \in Z$. This means that f is constant. □

Let $H \leq G$ be a subgroup of finite index, and fix some generating μ on G . It is well known that the subgroup H is μ -recurrent, namely, that the μ -random walk will hit H with probability 1. We denote the distribution of the first hit by θ . So θ is a generating measure on H which is called *the μ -hitting measure*. The stopping time theorem implies that the restriction map gives a canonical H -equivariant isomorphism between bounded μ -harmonic functions on G and bounded θ -harmonic functions on H . It follows that $\Pi(G, \mu)$ and $\Pi(H, \theta)$ are isomorphic as H -spaces.

Lemma 3.3. *Let G be a countable discrete group, μ be a generating probability measure on G , and let $g \in G$ have finite conjugacy class. Then g acts trivially on $\Pi(G, \mu)$.*

Proof. Let $g \in G$ be an element with a finite conjugacy class. Then its centralizer $H = C_G(g)$ is of finite index in G . Denote by θ the hitting measure on H . So $\Pi(G, \mu)$ and $\Pi(H, \theta)$ are H -isomorphic. Since g is in the center of H , by Theorem 3.2, it acts trivially on $\Pi(H, \theta)$, and hence it acts trivially on $\Pi(G, \mu)$. □

In particular, if there exists some generating measure μ such that the G -action on $\Pi(G, \mu)$ is faithful then G is an ICC group.

Proof of Proposition 3.1. Assume that there exists some μ on G for which $\Pi(G, \mu)$ is non-trivial. Let N denote the kernel of the action. So

G/N acts faithfully on $\Pi(G/N, \bar{\mu})$ where $\bar{\mu}$ is the μ -projected measure on G/N via $G \rightarrow G/N$. By Lemma 3.3, G/N is an ICC group. \square

Remark 3.4. *More generally, by applying an induction argument and Lemma 3.3, one can show that the hyper-FC-center [6, 28] acts trivially on $\Pi(G, \mu)$ for every generating μ on G .*

REFERENCES

- [1] Uri Bader and Yehuda Shalom, *Factor and normal subgroup theorems for lattices in products of groups*, Invent. Math. **163** (2006), no. 2, 415–454.
- [2] Laurent Bartholdi and Anna Erschler, *Poisson–Furstenberg boundary and growth of groups*, Probability Theory and Related Fields **168** (2017), no. 1-2, 347–372.
- [3] David Blackwell, *On transient markov processes with a countable number of states and stationary transition probabilities*, The Annals of Mathematical Statistics (1955), 654–658.
- [4] Nicolás Matte Bon, *Subshifts with slow complexity and simple groups with the liouville property*, Geometric and Functional Analysis **24** (2014), no. 5, 1637–1659.
- [5] Gustave Choquet and Jacques Deny, *Sur l'équation de convolution $\mu = \mu * \sigma$* , C. R. Acad. Sci. Paris **250** (1960), 799–801.
- [6] A Duguid, *A class of hyper-FC-groups*, Pacific Journal of Mathematics **10** (1960), no. 1, 117–120.
- [7] AM Duguid and DH McLain, *FC-nilpotent and FC-soluble groups*, Mathematical proceedings of the cambridge philosophical society, 1956, pp. 391–398.
- [8] Evgenii B Dynkin and MB Maljutov, *Random walk on groups with a finite number of generators*, Dokl. akad. nauk sssr, 1961, pp. 1042–1045.
- [9] Anna Erschler, *Boundary behavior for groups of subexponential growth*, Annals of Mathematics (2004), 1183–1210.
- [10] ———, *Liouville property for groups and manifolds*, Inventiones mathematicae **155** (2004), no. 1, 55–80.
- [11] Joshua Frisch and Pooya Vahidi Ferdowsi, *Non-virtually nilpotent groups have infinite conjugacy class quotients*, arXiv preprint arXiv:1803.05064 (2018).
- [12] Joshua Frisch, Omer Tamuz, and Pooya Vahidi Ferdowsi, *Strong amenability and the infinite conjugacy class property*, arXiv preprint arXiv:1801.04024 (2018).
- [13] Alex Furman, *Random walks on groups and random transformations*, Handbook of dynamical systems, 2002, pp. 931–1014.
- [14] H. Furstenberg and E. Glasner, *Stationary dynamical systems*, Contemp. Math., vol. 532, Amer. Math. Soc., Providence, RI, 2010.
- [15] Harry Furstenberg, *Noncommuting random products*, Transactions of the American Mathematical Society **108** (1963), no. 3, 377–428.
- [16] ———, *Random walks and discrete subgroups of Lie groups*, Advances in Probability and Related Topics **1** (1971), 1–63.
- [17] ———, *Boundary theory and stochastic processes on homogeneous spaces*, Harmonic analysis on homogeneous spaces **26** (1973), 193–229.
- [18] Shmuel Glasner, *On Choquet-Deny measures*, Ann. Inst. H. Poincaré B **12** (1976), 1–10.

- [19] ———, *Proximal flows*, Lecture Notes in Mathematics, Vol. 517, Springer-Verlag, Berlin-New York, 1976.
- [20] Yves Guivarc'h, *Croissance polynomiale et périodes des fonctions harmoniques*, Bull. Soc. Math. France **101** (1973), no. 333, 379.
- [21] Wojciech Jaworski, *Countable amenable identity excluding groups*, Canadian Mathematical Bulletin **47** (2004), no. 2, 215–228.
- [22] Wojciech Jaworski and C Robinson Edward Raja, *The Choquet–Deny theorem and distal properties of totally disconnected locally compact groups of polynomial growth*, New York J. Math **13** (2007), 159–174.
- [23] Vadim Kaimanovich, *The poisson boundary of covering markov operators*, Israel Journal of Mathematics **89** (1995), no. 1-3, 77–134.
- [24] Vadim A Kaimanovich, *Examples of non-abelian discrete groups with non-trivial exit boundary*, Zapiski Nauchnykh Seminarov POMI **123** (1983), 167–184.
- [25] ———, *Measure-theoretic boundaries of Markov chains, 0-2 laws and entropy*, Proceedings of the conference on harmonic analysis and discrete potential theory, 1992, pp. 145–180.
- [26] Vadim A Kaimanovich and Anatoly M Vershik, *Random walks on discrete groups: boundary and entropy*, The annals of probability (1983), 457–490.
- [27] Gregory A. Margulis, *Positive harmonic functions on nilpotent groups*, Soviet Math. Dokl. **7** (1966), 241–244.
- [28] DH McLain, *Remarks on the upper central series of a group*, Glasgow Mathematical Journal **3** (1956), no. 1, 38–44.
- [29] Joseph Rosenblatt, *Ergodic and mixing random walks on locally compact groups*, Mathematische Annalen **257** (1981), no. 1, 31–42.

(J. Frisch, O. Tamuz, P. Vahidi Ferdowsi) CALIFORNIA INSTITUTE OF TECHNOLOGY

(Y. Hartman) NORTHWESTERN UNIVERSITY